Guidebook 56th Annual Field Conference of Pennsylvania Geologists

Geology in the South Mountain Area, Pennsylvania



Hosts: September 26, 27, and 28, 1991 Dickinson College Carlisle, Pa. Pennsylvania Geological Survey

Guidebook for the

56th ANNUAL FIELD CONFERENCE OF PENNSYLVANIA GEOLOGISTS

GEOLOGY IN THE SOUTH MOUNTAIN AREA, PENNSYLVANIA

W. D. Sevon and Noel Potter, Jr., editors Contributors and field trip leaders: Rolf, Ackermann John H. Barnes Albert E. Becher S. W. Berkheiser, Jr. Douglas C. Chichester G. Michael Clark Gretchen Dockter Henry W. A. Hanson Marcus M. Key, Jr. Steven Lev Laura Pezzoli Robert J Schott Samuel J. Sims Samuel I. Root R. C. Smith, II Thomas Troy Randal L. Van Scyoc John H. Way Candie C. Wilderman Ellis L. Yochelson September 26-28, 1991 Hosts: Dickinson College Pennsylvania Geological Survey Headquarters Motel: The Embers Convention Center, Carlisle, PA Cover: Topographic maps at 1 inch = 1,600 feet for 3 stops on the field trip. Older maps are from a series with a contour interval of 10 feet that cover the entire South Mountain area in Pennsylvania. They were surveyed and drafted by Ambrose E. Lehman 1875-1885, and appear in Atlases D5 and D6 of the Second Pennsylvania Geological Survey. Modern maps are: Stop 3, USGS Iron Springs 7.5-minute quadrangle, C.I. = 20 feet. Stop 6, USGS Scotland and Caledonia Park 7.5-minute quadrangle, C.I. = 20 feet. Stop 8, USGS Dickinson and Mount Holly Springs 7.5-minute quadrangle, C.I. = 10 feet. Guidebooks distributed by: Field Conference of Pennsylvania Geologists, Inc. P. O. Box 1124 Harrisburg, PA 17108

0.9	50.0	Road on right leads to Kings Gap Environmental Educa-
		tion Center, a place well worth visiting.
3.1	52.1	Cross railroad. View to right of Stop 11.
0.9	54.0	TURN RIGHT onto Mountain View Road at Pennsy Supply
		sign. Traverse up a gravel surface.
0.8	54.8	SHARP TURN. BEAR LEFT.
0.1	54.9	TURN RIGHT following Mountain View Road.
0.2	55.1	GATE. Entrance to Pennsy Supply quarry.

STOP 11. GEOLOGY OF THE PENNSY SUPPLY QUARRY, MT. HOLLY, PENNSYLVANIA

Discussants: Marcus M. Key, Jr. and Samuel J. Sims.

Directions. Go through the gate and straight past the metal building on the right and up the hill. The road swings around to the right and heads up hill. At the fork, take the outside (right) lane which leads to Bench 1. Come down the same way.

GEOLOGY

The Pennsy Supply Quarry (formerly the R. A. Bender and Son Quarry) is 2.5 km west-south-west of Mt. Holly Springs in Dickinson Township, Cumberland County, Pennsylvania (Figure 75). The



Figure 75. Map showing location of quarry.

quarry is on the northwest slope of Mt. Holly at the northeast end of the Blue Ridge physiographic province between the Great Valley to the north and west and the Piedmont to the south and east. The quarry exploits the Antietam Quartzite, which was deposited during the Lower Cambrian (See Key, this guidebook). The Antietam Quartzite was later folded during the Paleozoic Appalachian orogenies that formed the South Mountain Anticlinorium. The quarry straddles the axis of the Bender's Quarry Anticline (Freedman, 1967; 1968) which plunges gently to the northeast. The present quarry is in the northwestern limb of this anticline (Sims, 1985).

The Antietam Quartzite lies on the upper phyllitic member of the Harpers Formation and is overlain by the Tomstown Dolomite. The Harpers Formation crops out upslope (to the south and east) from the quarry while the more-soluble and easily-eroded Tomstown Dolomite forms the southern margin of the Cumberland Valley to the north and west. The Harpers Formation and Tomstown Dolomite do not crop out in the quarry. Based on the distribution of float and topographic slope changes, the Antietam Quartzite is estimated to be at least 440 feet thick in Mt. Holly (Freedman, 1967).

The lithology of the Antietam Quartzite that is exposed in the quarry varies greatly. Textures range from fine- to coarsegrained, subangular to subrounded, well- to poorly-sorted, thinbedded to massive, with local crossbedding, laminations, and numerous *Skolithos linearis* worm tubes. Pre-deformation lithologies varied from thin clays, to quartz arenites, to granular quartz sandstones. Post-depositional deformation has altered these rocks to phyllites and quartzites. In some areas, slight schistose development is evident. The most common lithology is a relatively clean, massive, well-sorted, silica-cemented quartzite with up to 99 percent rounded quartz grains (Freedman, 1967; Fauth, 1968).

The Skolithos linearis worm tubes are ubiquitous in the Antietam Quartzite and are well illustrated in the slightly weathered, clean, indurated quartzites visible in the large blocks that have been set aside along the southwest edge of Bench 1 (Figure 76). Originally, these worm tubes had circular transverse, cross-sectional shapes. Because of subsequent stresses during the Appalachian orogenies, they have been deformed to a more-elliptical shape. Measurements of the resulting strain on the cross-sectional shapes of four worm tubes reveals a ratio of the short to long axis of 0.61.

The lithology and structure of the Antietam Quartzite are difficult to see in the quarry because of the massive nature of some units and well-developed jointing and weathering. It is especially difficult to determine the orientation of bedding. Look at the cliff between Benches 1 and 2 (Figures 76 and 77) for bedding planes. There are two reliable indicators of bedding plane surfaces: (1) the presence of clay layers interbedded in the quartzites and (2) *Skolithos linearis* worm tubes perpendicular to bedding. The strike of bedding ranges from N60°E to N90°E, and dips range from 45°NW to vertical to 30°SE; the dominant jointing strikes N47°E and dips 38°SE (Figure 77). These joints have been



Figure 76. Diagrammatic plan view of active portion of quarry as it appeared in June, 1991.

interpreted as extension joints parallel to the regional trend of the Bender's Quarry Anticline (Freedman, 1967). The divergent attitude of the joints and bedding results in blocky exposures.

Weathering causes the Antietam Quartzite to become extremely friable with a yellowish-tan to pinkish-red color. In some places such as in the fault zone along the crest of the anticlinal ridge, weathering may extend down as deep as 40 to 50 feet (Freedman, 1968).

What caused this deep weathering? Freedman (1967, 1968) argued that it was caused by leaching of the silica cement. He hypothesized that the silica dissolution was due to percolating groundwater which became alkaline as it seeped down through the overlying Tomstown Dolomite (Freedman, 1967). This does not seem plausible due to the difficulty of leaching silica in non-tropical climates (Friedman and Sanders, 1978).

alternative is that the deeply weathered zones re-Another flect a greater abundance of clay. To test this hypothesis, the type and relative abundance of clays in three Antietam Quartzite from the quarry were determined using X-ray diffraction. samples Results indicate that illite and kaolinite clays are present. The amount of illite is greatest in the clay layers interbedded between the quartzites, intermediate in the indurated quartzite, least in the friable quartzite. The amount of kaolinite is and greatest in the clay layers interbedded between the quartzites, intermediate in the friable quartzite, and least in the indurated



Figure 77. Diagrammatic cross section of Antietam Quartzite as it appeared in June, 1991 viewed from Bench 1 looking looking down strike (N77^OE).

quartzite. The interbedded clay layers are more commonly associated with the friable quartzite than the indurated quartzite. The deep weathering in some zones may simply indicate a greater kaolinite clay content. These deeply weathered zones are well known to the quarry operators because the clay-rich layers present a considerable problem of generation and disposal of slimes and are thus less desirable from an economic standpoint.

Spectacular liesegang rings are visible in the cliff between Benches 1 and 2 (Figures 76 and 77) as well as in the oversize blocks that have been set aside along the southwest edge of Bench 1 (Figure 76). Liesegang rings are secondary concentric rings formed by repeated precipitation within a fluid-saturated rock. Water in the once-saturated rock presumably transported iron in solution and precipitated it as hydrated iron oxide which we see today as rings of limonite. The liesegang rings found in the quarry are commonly formed around fractures in the quartzite. In the quarry, limonite can occasionally be found as pseudomorphs after another mineral, possibly pyrite. More commonly it occurs in botryoidal encrustations along fracture surfaces.

A problem is where did the iron come from because it is very rare in the Chilhowee Group. It may have come from dissolution of the overlying Tomstown Dolomite (Freedman, 1967). Dissolution

of the Tomstown Dolomite has been associated with the economic iron deposits on the south slope of South Mountain (Stose, 1932). diffraction analysis of three Antietam Quartzite samples X-ray from the quarry revealed the presence of some dolomite or high magnesium calcite which may indicate dissolution of the overlying Tomstown Dolomite and reprecipitation in the Antietam Quartzite. This hypothesis implies that the leisegang rings formed when the Tomstown Dolomite overlay the Antietam Quartzite. How long ago did the Tomstown Dolomite rest on the Antietam Quartzite at South Mountain? The fact that Triassic rocks rest on Cambrian rocks in the Gettysburg Basin, suggests that the erosion of the lower Paleozoic carbonates in that area occurred before or while the Triassic rocks were deposited. If this is also true for the South Mountain area, then the introduction of the iron from the Tomstown Dolomite into the Antietam Quartzite had to occur before or while the Triassic rocks were deposited. This would mean that the liesegang rings are at least Triassic in age.

Another line of reasoning suggests that the introduction of the iron may have been associated with the deep weathering of the Antietam Quartzite during the Middle Tertiary of later (See Clark, this guidebook). Could there have been Tomstown Dolomite overlying the Antietam Quartzite in the Middle Tertiary? Erosion for this area shed some light on the question. rates Sevon (1989)calculated the present erosion rate for the Juniata River drainage basin as 27 m/Ma based on suspended and dissolved loads in the river. From previous studies, Sevon (1989) compiled a erosion rate for limestone in the eastern United States of mean Using a value of 30 m/Ma and assuming erosion rates 32 m/Ma. have not changed since the Middle Tertiary (15 Ma), the net amount of denudation since this time is roughly 450 m. If we project the Tomstown up 450 m, it would overlie the Antietam where the quarry is currently located. In fact the current elevation difference between the Tomstown in the Cumberland Valley and the Antietam at the top of the quarry is only 230 m. The problem with this kind of exercise is that we know that erosion rates changed radically in the past 15 Ma as the South Mountain area was exposed to glacial and interglacial climates (Braun, 1989). This casts doubt on our ability to predict the past position of the Tomstown Dolomite relative to the Antietam Quartzite, but leaves open the possibility that its removal is a relatively recent event.

Are there other sources for the iron other than the Tomstown Dolomite? The Antietam Quartzite in the Mt. Holly area contains iron-rich heavy minerals such as magnetite (Freedman, 1967). The weathering of these may have been the source of iron in the liesegang rings. The amount of iron-rich heavy minerals in the Antietam Quartzite at Mt. Holly is unknown. This may be relatively unimportant because very little iron is required to stain a rock with liesegang rings (Pettijohn, 1975). The liesegang rings may have formed in association with the deep weathering in the Antietam Quartzite itself. This may have occurred long after the Tomstown eroded away.

The quarry is owned and operated by Pennsy Supply, Inc. of The quarry produces sand, washed-sand, and fine-Harrisburg. aggregate products for commercial use. The primary product is a concrete sand; other products are masonry sand, aggregate for concrete and asphalt, and general purpose sand and aggregate. There are two plants at the site for producing the range of products. The original plant that is nearest to the entrance to the quarry (Figure 75) has been upgraded and produces mainly masonry sand and sand approved by the Department of Environmental Resources (DER) for sand mounds for septic systems. This plant is dry plant and consists of a primary jaw crusher, a secondary a gyratory crusher, two sets of triple deck screens at 1 1/2 in., 3/8 in., and 5 mesh (4mm), a BARMAC impact crusher, and storage bins for loading directly into trucks or onto stackers for storage in stockpiles. The screens can be heated so that operation can continue during winter months.

The new and larger plant is both a dry and a wet plant sited to the west of the original plant (Figure 75). The dry part of the new plant consists of a primary jaw crusher, a secondary gyratory crusher, a BARMAC impact crusher, and triple deck screens set at 1 1/2, 1/2, and 3/8 inches. Material at minus 3/8 from the dry plant screens is fed to the wet plant that in. contains a triple deck wet screen set at 1/2, 1/4, and 3/16 in., an out-of-specification sand classifying tank, two screw classifiers, a cyclone, and a JADAIR clarifier. The main product from the wet plant is clean concrete sand, washed masonry sand, and clean fine Type A aggregate for asphalt that has a SRL rating of "E". The wet plant can be bypassed during cold weather, if necessary, to produce DER septic system sand, general purpose sand, Type A aggregate. Products are conveyed to stockpiles by and stackers around the plant and are loaded into trucks with front-end loaders.

The wet plant uses about 1,600 gallons of water per minute, which is supplied from a well that produces only about 100 gallons per hour. A well-designed recycling and storage program for plant water conserves almost all of the wash water. The JADAIR clarifier uses a flocculent to settle out the fine material in suspension that is fed from the classifier, and the clear water is recycled back through the plant. The flocculent is a polymer that neutralizes the naturally repelling cationic charges of the fine particles, which are then attracted to each other and flocculate and settle out of suspension (J. Kriz, personal communication, 1991). The slimes thus created are pumped from the clarifier as a slurry to settling ponds.

The quarry produces about 400,000 tons of product per year, about two-thirds of which is concrete sand. About 15 to 20 percent of the material quarried is lost as slimes and clay.

REFERENCES CITED

Adams, G. F., ed., 1975, Planation surfaces. peneplains, pediplains, and etchplains: Harrisburg, Dowden, Hutchinson and Ross, Inc., 476 p.

Aleinikoff, J. N., Zartman, R. E., Rankin, D. W., Lyttle, P. T., Burton, W. C., and McDowell, R. C., 1991, New U-Pb zircon ages for rhyolite of the Catoctin and Mount Rogers Formations-More evidence for two pulses of lapetan rifting in the central and southern Appalachians (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 2.

Allen, J. R. L., 1982, Sedimentary structures, their character and physical basis: New York, Elsevier Publishing Co., 1256 p.

Amsden, T. W. 1951, Paleontology of Washington County: Maryland Department of Geology and Mines, Water Resources Bulletin, No. 10, p. 98-123.

Anonymous, We. R., 1985, Gross Minerals, a force in the fine phyllite fillers field, Pennsylvania Geology, v. 16, no. 6, p. 2-5.

Badger, R. L. and Sinha, A. K., 1988, Age and Sr isotopic signature of the Catoctin volcanic province: Implications for subcrustal mantle evolution: Geology, v. 16, p. 692-695.

Badger, R. L. and Sinha, A. K., 1991, Nature of late Precambrian mafic magmatism within the Appalachian orogen (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 5.

Barrell, J., 1925, The nature and environment of the Lower Cambrian sediment of the southern Appalachians: American Journal of Science, v. 9, p. 1-20.

Barron, E. J., 1989, Climate variations and the Appalachians from the Late Paleozoic to the Present: results from model simulations, in Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 99-118.

Bascom, F., 1893, The structures, origin, and nomenclature of the acid volcanic rocks of South Mountain, Pennsylvania: Journal of Geology, v. 1, p. 813-832.

Bassler, R. S., 1919, Cambrian and Ordovician: Maryland Geological Survey, Johns Hopkins University Press, Baltimore, 408 p.

Baulig, H., 1952, Surfaces d'Aplanissement: Annales de Géographie, v. 61, p. 161-183; 245-262.

Becher, A. E. and Root, S. I., 1981, Groundwater and geology of the Cumberland Valley, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Water Resource Report 50, 95 p.

Becher, A. E. and Taylor, L. E., 1982, Groundwater resources in the Cumberland and contiguous valleys of Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Water Resource Report W-53, 67 p.

Berg, T. M., 1975, Geology and mineral resources of the Brodheadsville quadrangle, Monroe and Carbon Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 205a, 60 p.

Berg, T. M., and others, 1980, Geologic Map of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Map 1, scale 1:250,000.

Berg, T. M., Sevon, W. D., and Abel, R., 1984, Rock types of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Map 63, scale 1:500,000.

Berkheiser, S. W., Jr., 1985, High-purity silica occurrences in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 88, 67 p.

Berkheiser, S. W., Jr., Barnes, J. H., and Smith, R. C., II, 1985, Directory of the nonfuel-mineral producers in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Information Circular 54, 4th ed., 165 p.

Berkheiser, S. W., Jr., Potter, N., Jr., and Sevon, W. D., 1982, Hempt Brothers clay pit at Toland: in Potter, N., Jr., ed., South Mountain: Guidebook, Harrisburg Area Geological Society, p. 10-14.

Berkheiser, S. W., Jr. and Smith, R. C., II, 1990, Overhead protection, Pennsylvania's mineral industry contribution: Pennsylvania Geology, v. 21, no. 3, p. 2-6.

Bining, A. C., 1979, Pennsylvania iron manufacture in the eighteenth century, 2nd ed.: Harrisburg, Pennsylvania Historical and Museum Commission, 215 p.

Bjornerud, M. G., 1991, Conveying principles of finite strain with standard graphics software: Journal of Geological Education, v. 39, p. 23-27.

Black, R. F., 1976a, Periglacial features indicative of permafrost: ice and soil wedges: Quaternary Research, v. 6, p. 3-26.

Black, R. F., 1976b, Features indicative of permafrost: Annual Review of Earth and Planetary Sciences, v. 4, p. 75-94.

Black, R. F., 1983, Pseudo-ice-wedge casts of Connecticut, northeastern United States: Quaternary Research, v. 20, p. 74-89.

Blatt, H., 1980, Origin of sedimentary rocks, 2nd ed.: Englewood Cliffs, NJ, Prentice Hall, 782 p.

Bloomer, R. O., 1950, Late Pre-Cambrian and Lower Cambrian formations in central Virginia: American Journal of Science, v. 248, p. 753-783.

Bloomer, R. O. and Bloomer, R. R., 1947, The Catoctin Formation in central Virginia: Journal of Geology, v. 55, p. 94-106.

Bloomer, R. O. and Werner, H. J., 1955, Geology of the Blue Ridge in central Virginia: Geological Society of America Bulletin, v. 65, p. 579-606. Bond, G. C., Nickeson, P. A., and Kominz, M. A., 1984, Breakup of a supercontinent between 625 Ma and 555 Ma: New evidence and implications for continental histories: Earth and Planetary Science Letters, v. 70, p. 325-345.

Bowring, C. and Spencer, E., 1987, Catoctin pillow lavas (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 2, p. 77.

Bradt, P. T. and Wimpfheimer, L., 1987, Survey of Pocono lakes to evaluate sensitivity to acidification: Environmental Studies Center, Lehigh University, Bethlehem, PA.

Braun, D. D., 1989a, A revised Pleistocene glaciation sequence in eastern Pennsylvania: Support for limited Early Wisconsin ice and a single Late Illinoian advance beyond the Late Wisconsin border (abs.): 28th International Geological Congress Abstracts, v. 1: p. 1-196-1-197.

Braun, D. D., 1989b, Glacial and periglacial erosion of the Appalachians, <u>in</u> Gardner, T. W. and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 233-256.

Braun, D. D., and Inners, J. D., 1990, Weathering of conglomerate ledges and tors within the glacial limit in northeastern Pennsylvania: Evidence for single stage tor development under Pleistocene periglacial conditions (abs.): Geological Society of America Abstracts with Programs, v. 22, no. 2, p. 6.

Braun, E. L., 1950, Deciduous Forests of Eastern North America: Philadelphia, Blackiston Company, 596 p. Brezinski, D. K., 1991, Lithostratigraphy of the Lower Cambrian Tomstown Formation (abs.): Geological

Society of America Abstracts with Programs, v. 23, no. 1, p. 10.

Brown, W. R., 1970, Investigations of the sedimentary record in the Piedmont and Blue Ridge of Virginia, in Fischer, G. W., Pettijohn, F. J., Reed, J. C., Jr., and Weaver, K. N., eds., Studies of Appalachian Geology: Central and Southern: New York, Interscience, p. 335-349.

Bryan, K., Cleaves, A. B., and Smith, H. T. U., 1932/33, The present status of the Appalachian problem: Zeitschrift für Geomorphologie, v. 7, p. 312-320.

Bryan, K., 1946, Cryopedology the study of frozen ground and intensive frost-action with suggestions on nomenclature: American Journal of Science, v. 244, p. 622-642.

Büdel, J., 1982, Climatic Geomorphology: Princeton, Princeton University Press, 443 p.

Camuto, C., 1991, Dropping acid in the Southern Appalachians: A wild trout resource at considerable risk: Trout, Winter, 1991, p. 16-39.

Carter, B. J., and Ciolkosz, E. J., 1980, Soil temperature regimes of the Central Appalachians: Soil Science Society of America Journal, v. 44, p. 1052-1058.

Carter, B. J., and Ciolkosz, E. J., 1986, Sorting and thickness of waste mantle material in a sandstone spur in central Pennsylvania: Catena, v. 13, p. 241-256.

Cecil, C. B., 1990, Paleoclimatic controls on stratigraphic repetition of chemical and siliclastic rocks: Geology, v. 19, p. 533-536.

Ciolkosz, E. J., Carter, B. J., Hoover, M. T., Cronce, R. C., Waltman, W. J., and Dobos, R. R., 1990, Genesis of soils and landscapes in the Ridge and Valley province of central Pennsylvania, <u>in</u> Knuepfer, P. L. K. and McFadden, L. D., eds., Soils and landscape evolution: Amsterdam, Elsevier, p. 245-261.

Ciolkosz, E. J., Cronce, R. C., and Sevon, W. D., 1986, Periglacial features in Pennsylvania: University Park, PA, The Pennsylvania State University, Agronomy Department, Agronomy Series Number 92, 15 p.

Clark, G. M., 1989a, Central and southern Appalachian accordant ridge crest elevations south of glacial border: Regional erosional remnants or incipient periglacial surfaces? (abs.): 28th International Geological Congress Abstracts, v. 1, p. 1-299-1-300.

Clark, G. M., 1989b, Central and southern Appalachian water and wind gap origins: Review and new data, <u>in</u> Gardner T. W. and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 209-232.

Clark, G. M., and Ciolkosz, E. J., 1988, Periglacial geomorphology of the Appalachian Highlands and Interior Highlands south of the glacial border: Geomorphology, v. 1, p. 191-220.

- Clark, G. M., Ciolkosz, E. J., Kite, J. S., and Lietzke, D. A., 1989, Central and Southern Appalachian Geomorphology - Tennessee, Virginia, and West Virginia: 28th International Geological Congress Field Trip Guidebook T150, 105 p.
- Clark, G. M., and Hedges, J., in press, Origin of certain high-elevation local broad uplands in the Central Appalachians south of the glacial border - a paleoperiglacial hypothesis: 21st Annual Binghamton Geomorphology Symposium, Chichester, Wiley, in press.

Clark, M. J., ed., 1988, Advances in periglacial geomorphology: New York, John Wiley and Sons, 481 p.

Clarke, M. S., 1968, Pioneer iron works: Philadelphia, Chilton Book Company, 80 p.

Cleaves, E. T., Edwards, J. E., Jr., and Glaser, J. D., 1968, Geologic map of Maryland: Maryland Geological Survey, scale 1:250,000.

Cloos, E., 1947, Oolite deformation in the South Mountain fold, Maryland: Geological Society of America Bulletin, v. 58, p. 843-918.

Cloos, E., 1951, Physical features of Washington County: Maryland Department of Geology and Mines, Water Resources Bulletin 10, p. 17-94.

Cloos, E., 1958, Structural geology of South Mountain and Appalachians in Maryland: 23rd Annual Field Conference of Pennsylvania Geologists, Hagerstown, MD, Guidebook, 85 p. Cloos, E. and others, 1968, The geology of mineral deposits in south-central Pennsylvania: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, 80 p.

Cloos, E., 1971, Microtectonics along the western edge of the Blue Ridge, Maryland and Virginia: Johns Hopkins University, Studies in Geology No. 20, 234 p.

Cosby, B. J., Hornberger, G. M., Wright, R. F., and Galloway, J. N., 1986, Modeling the effects of acid deposition: control of long-term sulfate dynamics by soil sulfate adsorption: Water Resources Research, v. 22, p. 1283-1291.

Crimes, T. P., 1970, The significance of trace fossils in sedimentology, stratigraphy and palaeoecology with examples from Lower Palaeozoic strata, <u>in</u> Crimes, T. P. and Harper, J. C., eds., Trace Fossils: Geological Journal Special Issue 3, p. 106-126.

Daly, R. A., 1905, The accordance of summit levels among Alpine mountains: the fact and its significance: Journal of Geology, v. 13, p. 105-125.

Davis, W. N., 1889, The rivers and valleys of Pennsylvania: National Geographic Magazine, v. 1, p. 183-253.

Delcourt, P. A. and Delcourt, H. R., 1981, Vegetation maps for eastern North America; 40,000 yr BP to the present, <u>in</u> Romans, R. C., ed., Geobotany II: New York, Plenum, p. 123-165.

Delcourt, P. A. and Delcourt, H. R., 1984, Late Quaternary paleoclimates and biotic responses in eastern North America and the western North Atlantic Ocean: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 48: p. 263-284.

Demek, J., 1969, Cryoplanation terraces, their geographical distribution, genesis and development: Rozpravy Československé Akademie Věd. Rada Matematických A Přírodnich Věd, v. 79, no. 4, 80 p., 16 photos.

Demek, J., 1972, Manual of Detailed Geomorphological Mapping: Prague, Academia, 344 p.

d'Invilliers, E. V., 1887, Report on the iron ore mines and limestone quarries of Cumberland-Lebanon Valley, <u>in</u> Annual report of the Geological Survey for 1886: Pennsylvania Geological Survey, 2nd ser., Annual Report 1886, Pt. IV, p. 1411-1567.

Dockter, G. N., 1990, The Catoctin metabasalts: evidence of a rifting environment: unpublished independent research paper, Dickinson College, Carlisle, PA, 27 p.

Duffy, D. F. and Wittecar, G. R., 1991, Geomorphic development of segmented alluvial fans in the Shenandoah Valley, Stuarts Draft, Virginia (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 24.

Edwards, J., Jr., compiler, 1978, Geologic map of Washington County: Maryland Geological Survey, scale 1:62,500.

Ehlen, J., 1990, Geomorphic, petrographic and structural classification of granite landforms using spatial patterns (abs.): Geological Society of America Abstracts With Programs, v. 22, no. 7, p. A21.

Espenshade, G. H., 1986, Geology of the Marshall quadrangle, Faquier County, Virginia: U. S. Geological Survey Bulletin 1560, 59 p.

Faill, R. T. and MacLachlan, D. B., 1989, Tectonic terranes of southeastern Pennsylvania (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 1, p. 13.

Farlekas, G., 1961, The geology of part of South Mountain of the Blue Ridge province north of the Pennsylvania-Naryland border: unpublished M.S. thesis, University Park, The Pennsylvania State University, 64 p.

Fauth, J. L., 1968, Geology of the Caledonia Park quadrangle Area, South Mountain, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 129a, 133 p.

Fauth, J. L., 1977, Geologic map of the Catoctin Furnace and Blue Ridge Summit Quadrangles, Maryland: Maryland Geological Survey, scale 1:24,000.

Fauth, J. L., 1978, Geology and mineral resources of the Iron Springs area, Adams and Franklin Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 129c, 72 p.

Fauth, J. L., 1981, Geologic map of the Myersville Quadrangle, Maryland: Maryland Geological Survey, scale 1:24,000.

Fedonkin, M. A., 1981, Palaeoichnology of the Precambrian-Cambrian transition, in Taylor, M. E., ed., Short papers from the Second International Symposium on the Cambrian System: U.S. Geological Survey Open-File Report 81-743, p. 89-90.

Fenneman, N. M., 1938, Physiography of eastern United States: New York, McGraw-Hill Book Company, 714 p.

Flemal, R. C., 1971, The attack on the Davisian system of geomorphology: a symposium: Journal of Geologic Education, v. 19, p. 3-13.

Flinn, D. 1962, On folding during three-dimensional progressive deformation: Geological Society of London Quarterly Journal, v. 118, p. 385-433.

Flippo, H. N., Jr., 1974, Springs of Pennsylvania: Pennsylvania Department of Environmental Resources, Water Resources Bulletin 10, 46 p.

Flower, L. E., 1975, History of Pine Grove Furnace: Carlisle, PA, Cumberland County Historical Society, 21 p.

Foose, R. M., 1945a, Iron-manganese ore deposits of White Rocks, Cumberland County: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 26, 35 p.

Foose, R. M., 1945b, Manganese minerals of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 27, 130 p.

Foose, R. M. and others, 1948, Excursion 1, South Mountain, in Cathcart, S. H., ed., 14th Annual Field

Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 1-6.

Frakes, L. A., 1979, Climates throughout geologic time: Amsterdam, Elsevier Science Publishers, 310 p. Freedman, J., 1967, Geology of a portion of the Mount Holly Springs Quadrangle, Adams and Cumberland

Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Progress Report 169, 66 p. Freedman, J., 1968, Bender's Quarry - Nt. Holly Springs, Pennsylvania, <u>in</u> The geology of mineral deposits in

south-central Pennsylvania: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 28-37.

French, H. M., 1976, The periglacial environment: London, Longman, 309 p.

- French, H. M. and Karte, J., 1988, A periglacial overview, <u>in</u> Clark, M. J., ed., Advances in Periglacial Geomorphology: New York, John Wiley and Sons, p. 463-473.
- Friedman, G. M. and Sanders, J. E., 1978, Principles of Sedimentology: New York, John Wiley and Sons, 792 p. Fritz, W. H., 1972, Lower Cambrian trilobites from the Sekawi Formation type section, Mackenzie Mountains, north-western Canada: Geological Survey of Canada Bulletin, v. 212, p. 1-58.

Fritz, W. H. and Crimes, T. P., 1985, Lithology, trace fossils, and correlation of the Precambrian-Cambrian boundary beds, Cassiar Mountains, north-central British Columbia: Geological Survey of Canada Paper 83-13, p. 1-24.

Fulton, R. J., 1989, ed., Quaternary Geology of Canada and Greenland: Geological Society of America, Volume K-1, Geological Survey of Canada, no. 1, 839 p.

Gagen, C. J. and Sharpe, W. E., 1987, Net sodium loss and mortality of three salmonid species exposed to a stream acidified by atmospheric deposition: Environmental Contamination and Toxicology Bulletin, v. 39, p. 7-14.

Gannett Flemming Corddry and Carpenter, Inc., 1945, Report on water treatment project for the Borough of Chambersburg, PA: Harrisburg, PA.

Gardiner, V. and Dackombe, R., 1983, Geomorphological Field Manual: London, George Allen and Unwin, 254 p. Gardner, T. W. and Sevon, W. D., eds., 1989, Appalachian Geomorphology: Amsterdam, Elsevier, 318 p.

Geiser, P. A. and Engelder, T., 1983, The distribution of layer-parallel shortening fabrics in the Appalachian foreland of New York and Pennsylvania: Evidence for two non-coaxial phases of the Alleghanian orogeny, <u>in</u> Hatcher, R. D., Jr., Williams, H., and Zeitz, I., eds., Contributions to the

tectonics and geophysics of mountain chains: Geological Society of America Memoir 158, p. 161-175. Gerhart, J. M. and Lazorchick, G. J., 1988, Evaluation of the ground-water resources of the lower Susquehanna River Basin, Pennsylvania and Maryland: U.S. Geological Survey Water-Supply Paper 2284, 128 p.

Geyer, A. R., Smith, R. C., II, and Barnes, J. H., 1976, Mineral collecting in Pennsylvania: Pennsylvania Geological Survey, 4th ser., General Geology Report 33, 260 p.

Godfrey, A. E., 1975, Chemical and physical erosion in the South Mountain anticlinorium, Maryland: Maryland Geological Survey, Information Circular 19, 35 p.

Godfrey, A. E. and Cleaves, E. T., 1991, Landscape analysis: Theoretical considerations and practical needs: Environmental Geology and Water Sciences, v. 17, no. 2, p. 141-155.

Goldthwait, R. P., 1976, Frost sorted patterned ground: A review: Quaternary Research, v. 6, p. 27-35.

Gordon, S. E., 1922, The mineralogy of Pennsylvania: Philadelphioa, Academy of Natural Sciences Special Publication 1, 158 p.

Grimvall, A., Cole, C. A., Allard, B., and Sanden, P., 1986, Quality trends of public water supplies in Sweden: Water Quality Bulletin, v. 11, no. 4, p. 6-11.

Hack, J. T., 1960, Interpretation of erosional topography in humid temperate regions: American Journal of Science Bradley Volume, v. 258-A, p. 80-97.

Hack, J. T., 1965, Geomorphology of the Shenandoah Valley Virginia and West Virginia and origin of the residual ore deposits: U.S. Geological Survey Professional Paper 484, 84 p.

Hack, J. T., 1975, Dynamic equilibrium and landscape evolution, in Melhorn, W. N. and Flemal, R. C., eds., Theories of landform development: London, George Allen and Unwin, p. 87-102.

Haines, T. A., 1981, Acidic precipitation and its consequences for aquatic ecosystems: a review: Transactions of the American Fisheries Society, v. 110, p. 669-707.

Hatcher, R. D., Jr., 1990, Structural geology: principles, concepts, and problems: Columbus, OH, Merrill Publishing Co., 531 p.

Hatcher, R. D., Thomas, W. A., and Viele, G. W., 1989, The Appalachian-Ouachita Orogen in the United States: The Geology of North America, v. F-2: Geological Society of America, 767 p.

Hedges, J., 1969, Opferkessel: Zeitschrift für Geomorphologie, v. 13, p. 22-55.

Hedges, J., 1975, Multiple cycles of cryoplanation on Sugarloaf Mountain, Maryland: Biuletyn Peryglacjalny, v. 24, p. 233-243.

Heron, S. D., Jr., Judd, J. B., and Johnson, H. S., Jr., 1971, Clastic dikes associated with soil horizons in the North and South Carolina Coastal Plain: Geological Society of America Bulletin, v. 82, p. 1801-1810.

Hofmann, A. W., 1988, Chemical differentiation of the Earth: the relationship between mantle, continental crust, and oceanic crust: Earth and Planetary Science Letters, v. 90, p. 297-314.

Hosterman, J. W., 1968, White clay deposits near Mt. Holly Springs, Cumberland County, Pennsylvania, in The geology of mineral deposits in south-central Pennsylvania: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 38-51. Hosterman, J. W., 1969, White clay deposits near Mt. Holly Springs, Cumberland County, Pennsylvania: U.S. Geological Survey Professional Paper 650B, p. 66-72.

Hupp, C. R., 1983, Geo-botanical evidence of Late Quaternary mass wasting in block field areas of Virginia: Earth Surface Processes and Landforms, v. 8, p. 439-450.

Jacobson Jr., G. L., Webb III, T., and Grimm, E. C., 1987, Patterns and rates of vegetation change during the deglaciation of eastern North America, <u>in</u> Ruddiman W. F. and Wright, H. E., Jr., eds., North America and adjacent oceans during the last deglaciation: Geological Society of America, p. 277-288.

Johnson, F., 1986, Acid precipitation: Pennsylvania Fish Commission, PFC 1003 7/86 10 M.

Karrasch, H., 1974, Hangglättung und Kryoplanation an Beispielen aus den Alpen und kanadischen Rocky Mountains: Akademie Der Wissenschaften, Göttingen. Mathematische-Physikale Abhandlungen, v. 29, p. 287-300.

Karte, J., 1982, Development and present state of German periglacial research in arctic and alpine environments: Biuletyn Peryglacjalny, v. 29, p. 183-201.

Kauffman, M. E. and Frey, E. P., 1979, Antietam sandstone ridges - exhumed barrier islands or fault-bounded blocks? (abs.): Geological Society of America Abstracts with Programs, v. 11, no. 1, p. 18.

Kaufmann, P. R., Herlihy, A. T., Elwood, J. W., Mitch, M. E., Overton, W. S., Sale, M. J., Messer, J. J., Cougan, K. A., Peck, D. V., Reckhow, K. H., Kinney, A. J., Christie, S. J., Brown, D. D., Hagley, C. A., and Jager, H. I., 1988, Chemical characteristics of streams in the mid-Atlantic and southeastern United States (Results of the National Stream Survey Phase 1), in Vol. 1: Population descriptions and physico-chemical relationships: U.S. Environmental Protection agency, Washington, D.C., EPA/600/3-88/021a.

Kay, M., 1951, North American Geosynclines: Geological Society of America Memoir 48, 143 p.

Kemper, J., III, 1968?, American charcoal making: National Park Service, U.S. Department of the Interior, Eastern National Park and Monument Association, 25 p.

Kimmel, W. G., Murphy, D. J., Sharpe, W. E., and DeWalle, D. R., 1985, Macroinvertebrate community structure and detritus processing rates in two southwestern Pennsylvania streams acidified by atmospheric deposition: Hydrobiologia, v. 124, p. 97-102.

King, P. B., 1949, The base of the Cambrian in the southern Appalachians: American Journal of Science, v. 247, p. 513-530; 622-645.

King, P. B., 1950, Geology of the Elktin area, Virginia: U.S. Geological Survey Professional Paper 230, 82 p.

King, P. B. and Fergusen, H. W., 1960, Geology of northeasternmost Tennessee: U.S. Geological Survey Professional Paper 311, 136 p.

Kline, S. W., Conley, J. F., and Evans, N., 1987, The Catoctin Formation in the eastern Blue Ridge of Virginia: Evidence for submarine volcanism (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 2, p. 93.

Kochel, R. C., 1990, Humid fans of the Appalachian Mountains, <u>in</u> Rachocki, A. H. and Church, M., eds., Alluvial fans: New York, John Wiley and Sons, p. 109-129.

Kulander, B. R. and Dean, S. L., 1986, Structure and tectonics of the central and southern Appalachian Valley and Ridge Plateau Provinces, West Virginia, and Virginia: American Association of Petroleum Geologists Bulletin, v. 70, p. 1674-1684.

Kurjack, D. C., 1958, Hopewell Village national historic site: Washington, D.C., National Park Service Historical Handbook Series No. 8, 44 p.

Lachenbruch, A. H., 1962, Mechanics of thermal contraction cracks and ice-wedge polygons in permafrost: Geological Society of America Special Paper 70, 69 p.

Laurence, R. A. and Palmer, A. R., 1963, Age of the Murray Shale and Hesse Quartzite on Chilhowee Mountain, Blount County, Tennessee: U.S. Geological Survey Professional Paper 475C, p. 53-54.

Lauriol, B., 1990, Canadian landform examples 18: Cryoplanation terraces, northern Yukon: Le Geographe canadien, v. 34, no. 4, p. 347-351.

Leffingwell, E. de K., 1915, Ground-ice wedges. The dominant form of ground-ice on the north coast of Alaska: Journal of Geology, v. 23, p. 635-654.

Leffler, R. J., 1981, Estimating average temperatures on Appalachian summits: Journal of Applied Meteorology, v. 20, p. 637-642.

Lesley, J. P., 1892, A summary description of the geology of Pennsylvania: Pennsylvania Geological Survey, 2nd ser., v. 1, 719 p.

Liestøl, O., 1961, Talus taluses in Arctic regions: Norsk Polarinstitutt Årbok, p. 102-105.

Lisle, R. J., 1985, Geological strain analysis, a manual for the Rf/f technique: New York, Pergamon Press, 99 p.

Løzinski, M. W., 1909, Über die mechanische Verwitterung der Sandsteine im gemässigten Klima: Bulletin International Akademie Sciences Crakovie, Classe Scientific Mathematische Naturelles, v. 1, p. 1-25.

Lynch, J. A., Corbett, E. S., and Kostelnik, K. M., 1986, Atmospheric deposition: spatial and temporal variation in Pennsylvania, 1985: Institute for Research on Land and Water Resources, The Pennsylvania State University, University Park, PA.

Lynch, J. A., Corbett, E. S., and Kostelnik, K. N., 1987, Atmospheric deposition: spatial and temporal variation in Pennsylvania, 1986: Institute for Research on Land and Water Resources, The Pennsylvania

State University, University Park, PA.

M'Cauley, I. H., 1878, Historical sketch of Franklin County, Pennsylvania: Chambersburg, PA, D. F. Pursel, 322 p.

Mackay, J. R., 1974, Ice-wedge cracks, Garry Island, Northwest Territories: Canadian Journal of Earth Science, v. 11, p. 1366-1383.

Mackay, J. R., 1990, Some observations on the growth and deformation of epigenetic, syngenetic, and anti-syngenetic ice wedges: Permafrost and Periglacial Processes, v. 1, p. 15-29.

MacLachlan, D. B. and Root, S. I., 1966, Comparative tectonics of the Cumberland and Lebanon Valleys: 31st Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, 90 p.

McDonald, M. E., 1985, Acid deposition and drinking water: Environmental Science and Technology, v. 19, p. 772-776.

McDonald, M. G. and Harbaugh, A. W., 1988, A modular three-dimensional finite-difference ground-water flow model: U.S. Geological Survey Techniques of Water Resources Investigations, book 6, chap. A1.

Middlekauff, B. D., 1987, Relict periglacial morphosequences in the northern Blue Ridge: unpublished Ph.D. thesis, East Lansing, Michigan State University, 172 p.

Mills, H. H., 1981, Boulder deposits and the retreat of mountain slopes, or, "gully gravure" revisited: Journal of Geology, v. 89, p. 649-660.

Mills, H. H., Brakenridge, G. R., Jacobson, R. B., Newell, W. L., Pavich, M. J., and Pomeroy, J. S., 1987, Appalachian Mountains and Plateaus: Geological Society of America, Centennial Volume 2, p. 5-50.

Mills, H. H. and Delcourt, P. A., in press, Appalachian highlands and interior low plateaus, <u>in</u> Morrison, R. B., ed., Quaternary non-glacial geology of the conterminous United States: Geological Society of America, Volume K-2, in press.

Monmonier, M. S., 1967, Upland accordance in the Ridge and Valley section of Pennsylvania: unpublished M.S. thesis, University Park, The Pennsylvania State University, 58 p.

Monmonier, M. S., 1968, Trends in upland accordance in Pennsylvania's Ridge and Valley section: Pennsylvania Academy of Science Proceedings, v. 42, p. 157-162.

Monmonier, M. S., 1971, Upland adjustment to regional drainage in central Pennsylvania: an application of trend surface analysis: Journal of Geography, v. 70, p. 360-370.

- Mose, D. G., Diecchio, R. J., DiGuiseppi, W. H., and Nagel, M. S., 1985, Confirmation of a latest Precambrian (2600 m.y.) age for the Catoctin Formation in Virginia (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 2, p. 126.
- Moss, J. H., 1976, Periglacial origin of extensive lobate colluvial deposits on the south flank of Blue Mountain near Shartlesville and Strausstown, Berks County, Pennsylvania: Pennsylvania Academy of Science Proceedings, v. 50, p. 42-44.

Nagel, M. S. and Mose, D. G., 1984, A revised geochemical and geochronological picture of the Catoctin Formation (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 182.

National Atmospheric Deposition Program, 1990, NADP/NTN annual data summary, precipitaion chemistry in the United States, 1989.

Neal, J. T., Langer, A. M., and Kerr, P. F., 1968, Giant desiccation polygons of Great Basin playas: Geological Society of America Bulletin, v. 79, p. 69-90.

Nelson, A., 1968, GAF Corporation quarries--Charmian, Pennsylvania, <u>in</u> Cloos, E. and others, The geology of mineral deposits in south-central Pennsylvania, Annual Field Conference of Pennsylvania Geologists, 33rd, Harrisburg, PA, Guidebook, p. 6-10.

Nickelsen, R. P., 1956, Geology of the Blue Ridge near Harpers Ferry, West Virginia: Geological Society of America Bulletin, v. 67, p. 239-270.

Nickelsen, R. P., 1979, Sequence of structural stages of the Allegheny orogeny at the Bear Valley strip mine, Shamokin, Pennsylvania: American Journal of Science, v. 279, p. 225-271.

Nutter, L. J., 1973, Hydrogeology of the carbonate rocks, Frederick and Hagerstown Valleys Maryland: Maryland Geological Survey, Report of Investigations No. 19, 70 p.

Olson, C. G., 1989, Mountain soils of eastern Appalachians (abs.): 28th International Geological Congress Abstracts. v. 2, p. 2-548.

Palmer, A. R., 1981, Subdivision of the Sauk Sequence, in Taylor, M. E., ed., Short Papers from the Second International Symposium on the Cambrian System: U.S. Geological Survey Open-File Report 81-743, p. 160-161.

Palmer, A. R., 1983, The Decade of North American Geology, 1983 geologic time scale: Geology, v. 11, p. 503-504.

Patterson, J. G. and Simpson, E. L., 1991, Paleoenvironmental constraints on the Chilhowee Group of the eastern Blue Ridge: Thoroughfare Gap, Virginia (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 113.

Pavich, M. J., 1989, Regolith residence time and the concept of surface age of the Piedmont "peneplain", in Gardner, T. W. and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 181-196.

Pearce, J. A. and Cann, J. R., 1973, Tectonic setting of basic volcanic rocks determined using trace element analysis: Earth and Planetary Science Letters, v. 19, p. 290-300.

- Pearse, J. B., 1970, A concise history of the iron manufacture of the American colonies up to the Revolution and of Pennsylvania up to the present time: New York, Burt Franklin, Research and Source Works Series 497, Selected essays in history, economics, and social sciences 142 (originally published in 1876, author "metallurgist, engineer, Commissioner of Geological Survey of Pennsylvania").
- Peltier, L. C., 1949, Pleistocene terraces of the Susquehanna River, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Bulletin G 23, 158 p.
- Peltier, L. C., 1950, The geographic cycle in periglacial regions as it is related to climatic geomorphology: Association of American Geographers Annals, v. 40, p. 214-236.
- Pettijohn, F. J., 1957, Sedimentary rocks: New York, Harper and Row, 628 p.
- Pettyjohn, W. A. and Henning, R., 1979, Preliminary estimate of ground-water recharge rates, related streamflow and water quality in Ohio: The Ohio State University Department of Geology and Mineralogy Project Completion Report No. 522, 323 p.
- Péwé, T. L., 1970, Altiplanation terraces of early Quaternary age near Fairbanks, Alaska: Acta Geographica Lodziensia, v. 24, p. 357-363.
- Péwé, T. L., 1975, Quaternary geology of Alaska: U.S. Geological Survey Professional Paper 835, 145 p.
- Péwé, T. L., 1983, The periglacial environment in North America during Wisconsin time, <u>in</u> Porter, S. C., ed., Late-Quaternary Environments of the United States. Volume 1. The Late Pleistocene: Minneapolis, University of Minnesota Press, p. 157-189.
- Péwé, T. L., Church, R. E., Andresen, M. J., 1969, Origin and paleoclimatic significance of large-scale patterned ground in the Donnelly Dome area, Alaska: Geological Society of America Special Paper 103, 87 p.
- Pierce, K. L., 1965, Geomorphic significance of a Cretaceous deposit in the Great Valley of southern Pennsylvania: U.S. Geological Survey Professional Paper 525-C, p. C152-C156.
- Pierce, K. L., 1966, Bedrock and surficial geology of the McConnellsburg quadrangle, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 109a, 111 p.
- Poag, C. W., and Sevon, W. D., 1989, A record of Appalachian denudation in postrift Mesozoic and Cenozoic sedimentary deposits of the U.S. Middle Atlantic continental margin, <u>in</u> Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 119-157.
- Priesnitz, K., 1988, Cryoplanation, <u>in</u> Clark, M. J., ed., Advances in Periglacial Geomorphology: Chichester, John Wiley and Sons, p. 49-67.
- Ramsay, J. G., 1967, Folding and fracturing of rocks: New York, McGraw-Hill, 568 p.
- Ramsay, J. G. and Huber, M. I., 1983, The tectonics of modern structural geology, Vol. 1, Strain Analysis: New York, Academic Press, Inc., 307 p.
- Rankin, D. W., 1967, Guide to the geology of the Mount Rogers area, Virginia, North Carolina, and Tennessee: Carolina Geological Society Field Trip, Guidebook, 48 p.
- Rankin, D. W., 1975, The continental margin of eastern North America in the southern Appalachians: The opening and closing of the Proto-Atlantic Ocean: American Journal of Science, v. 275-A, p. 298-336.
- Rankin, D. W., Stern, T. W., Reed, J. C., Jr., and Newell, M. F., 1969, Zircon ages of felsic rocks in the Upper Precambrian of the Blue Ridge, Appalachian Mountains: Science, v. 166, p. 741-744.
- Reed, J. C., 1955, Catoctin Formation near Luray, Virginia: Geological Society of America Bulletin, v. 66, p. 871-896.
- Reger, R. D. and Péwé, T. L., 1976, Cryoplanation terraces: indicators of a permafrost environment: Quaternary Research, v. 6, p. 99-109.
- Resser, C. E. and Howell, B. J., 1938, Lower Cambrian Olenellus zone of the Appalachians: Geological Society of America Bulletin, v. 49, p. 195-248.
- Rogers, H. D., 1858, The geology of Pennsylvania: Philadelphia, v. 1, 586 p.
- Root, S. I., 1968, Geology and mineral resources of southeastern Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 119cd.
- Root, S. I., 1970, Structure of the northern terminus of the Blue Ridge in Pennsylvania: Geological Society of America Bulletin, v. 81, p. 815-830.
- Root, S. I., 1971, Geology and mineral resources of northeastern Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 119ab, 104 p.
- Root, S. I., 1977, Geology and mineral resources of the Harrisburg West area of Cumberland and York Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 148ab, 106 p.
- Root, S. I., 1978, Geology and mineral resources of the Carlisle and Mechanicsburg Quadrangles, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 138ab, 1 p.
- Root, S. I., 1989, Basement control of structure in the Gettysburg rift basin, Pennsylvania and Maryland: Tectonophysics, v. 166, p. 281-292.
- Root, S. I. and Hoskins, D. M., 1977, Lat. 40⁰N fault-zone, Pennsylvania: a new interpretation: Geology, v. 5, p. 719-723.
- Root, S. I. and MacLachlan, D. B., 1978, Western limit of Taconic allochthons in Pennsylvania: Geological Society of America Bulletin, v. 89, p. 1515-1528.
- Rorabaugh, M. I., 1964, Estimating changes in bank storage and ground-water contribution to streamflow:

International Association of Scientific Hydrology Publication no. 63, p. 432-441.

Rose, A. W., 1970, Metal mines and occurrences in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resources Report 50, Part 3, 14 p.

Russell, R. J., 1933, Alpine land forms in western United States: Geological Society of America Bulletin, v. 44, p. 927-950.

Saad, D. A. and Hippe, D. J., 1990, Large springs in the Valley and Ridge physiographic province of Pennsylvania: U.S. Geological Survey Open-File Report 90-164, 17 p.

Schmidlin, T. W., 1982, Leffler's method of estimating average temperatures of Appalachian summits: Evaluation in New York: Journal of Applied Meteorology, v. 21, p. 745-747.

Schofield, C. and Trojnar, J. R., 1980, Aluminum toxicity to brook trout (<u>Salvelinus fontinalis</u>) in acidified waters, <u>in</u> Toribara, T. Y., Miller, M. W., and Morrow, P. E., eds., Polluted Rain: New York, Plenum Press, p. 341-365.

Schwab, F. L., 1970, Origin of the Antietam Formation (Late Precambrian?-Lower Cambrian), central Virginia: Journal of Sedimentary Petrology, v. 40, p. 354-366.

Schwab, F. L., 1972, The Chilhowee Group and the Late Precambrian-Early Paleozoic sedimentary framework in the central and southern Appalachians, in Lessing, P. and others, eds., Appalachian structures - origin, evolution, and possible potential for new exploration frontiers: Morgantown, West Virginia University and West Virginia Geological and Economic Survey, p. 59-101.

Scotese, C. R. and others, 1979, Paleozoic base maps: Journal of Geology, v. 87, p. 217-277.

Sears, C. E., 1957, Late Cretaceous erosion surface in southwest Virginia (abs.): Geological Society of America Bulletin, v. 68, p. 1883.

Seilacher, A., 1967, Bathymetry of trace fossils: Marine Geology, v. 5, p. 413-428.

Sepkoski, J. J., Jr. and Knoll, A. H., 1983, Precambrian-Cambrian boundary: The spike is driven and the monolith crumbles: Paleobiology, v. 9, p. 199-206.

Sevon, W. D., 1984, A sandstone weathering rate from northeastern Pennsylvania (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 1, p. 63.

Sevon, W. D., 1985, Pennsylvania's Polygenetic Landscape: Harrisburg Area Geological Society, 4th Annual Field Trip Guidebook, 55 p.

Sevon, W. D., 1987, The Hickory Run boulder field, a periglacial relict, Carbon County, Pennsylvania, in Roy, D. C., ed., Northeastern Section of the Geological Society of America, Centennial Field Guide, Vol. 5: Geological Society of America DNAG Volume, p. 75-76.

Sevon, W. D., 1989, Erosion in the Juniata drainage basin, Pennsylvania, in Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 303-318.

Sevon, W. D., 1990, The Hickory Run boulder field, Hickory Run State Park, Carbon County, Pennsylvania: Northeastern Geology, v. 12, p. 42-45.

Sevon, W. D., Potter, Jr., N., and Crowl, G. H., 1983, Appalachian peneplains: an historical review: Earth Sciences History, v. 2, p. 156-164.

Sharpe, W. E., DeWalle, D. R., Liebfried, R. T., Dinicola, R. S., Kimmel, W. G., and Sherwin, L. S., 1984, Causes of acidification of four streams on Laurel Hill in southwestern Pennsylvania: Journal of Environmental Quality, v. 13, p. 619-631.

Shirk, W. R., 1980, Geology of southcentral Pennsylvania: Chambersburg, PA, Robson and Kaye, Inc., 136 p.

Siddens, L. K., Seim, W. K., and Curtis, L. R., 1986, Comparison of continuous and episodic exposure to acidic, aluminum-contaminated waters of brook trout (<u>Salvelinus fontinalis</u>): Canadian Journal of Fisheries and Aquatic Sciences, v. 43, p. 2036-2040.

Simpson, E. L., Linski, D., Mull, M. F., Keiser, J. P., Horsnall, S. L., and Hendricks, J. S., 1991, Depositional processes in outer-shelf sediments of the Lower Cambrian Harpers Formation of the Chilhowee Group, south-central Pennsylvania (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 127.

Simpson, E. L. and Sundberg, F. A., 1987, Early Cambrian age for synrift deposits of the Chilhowee Group of southwestern Virginia: Geology, v. 15, p. 123-126.

Simpson, K. W., Bode, R. W., and Colquhoun, J. R., 1985, The macroinvertebrate fauna of an acid-stressed headwater stream in the Adirondack Mountains, New York: Freshwater Biology, v. 15, p. 671-681.

Sims, S. J., 1985, Geological reconnaissance survey and reserve estimate - R. A. Bender & Son Sand Quarry, Mt. Holly Springs, Dickinson Township, Cumberland County, Pennsylvania: unpublished private report for Pennsy Supply, Inc., 18 p.

Sloss, L. L., 1963, Sequences in the cratonic interior of North America: Geological Society of America Bulletin, v. 74, p. 93-114.

Smith, H. T. U., and Smith, A. P., 1945, Periglacial rock streams in the Blue Ridge area (abs.): Geological Society of America Bulletin, v. 56, p. 1198.

Smith, R. C., II, 1978, The mineralogy of Pennsylvania, 1966-1975: Friends of Mineralogy, Pennsylvania Chapter, Special Publication No. 1, 304 p.

Smith, R. C., II and Berkheiser, S. W., Jr., 1985, Silica 99 and 44/100% pure, waiting to be be mined: Pennsylvania Geology, v. 16, no. 1, p. 2-4. Spencer, E. W., 1988, Introduction to the structure of the Earth: New York, McGraw-Hill, 551 p.

Spencer, E. W., Bell, J. D., and Kozak, S. J., 1989, Valley and Ridge and Blue Ridge traverse, central Virginia: 28th International Geological Congress, Washington, D.C., Field Trip Guidebook, <u>in</u> Glacial geology and geomorphology of North America, vol. 2: American Geophysical Union, 69 p.

Stone, R. W., 1923, Roofing granules industry in southeastern Pennsylvania: Pennsylvania Geological Survey, 4th ser., Progress Report 82, 4 p.

Stone, R. W. and Hickok, W. O., 1933, Trip 4, South Mountain, <u>in</u> Ashley, G. W., ed., 3rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 11-14.

Stose, G. W., 1906, The sedimentary rocks of South Mountain, Pennsylvania: Journal of Geology, v. 14, p. 201-220.

Stose, G. W., 1907, White clay of South Mountain, Pennsylvania: U.S. Geological Survey Bulletin, v. 315, p. 322-334.

Stose, G. W., 1909, Mercersburg-Chambersburg folio: U.S. Geological Survey Geologic Atlas, Folio No. 170, 19 p.

Stose, G. W., 1932, Geology and mineral resources of Adams County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Bulletin C1, 153 p.

Stose, G. W., 1953, Geology of the Carlisle Quadrangle, Pennsylvania: U.S. Geological Survey Geological Quadrangle Map GQ-28.

Stose, G. W. and Bascom, F., 1929, Fairfield-Gettysburg folio: U.S. Geological Survey Geologic Atlas, Folio No. 225, 22 p.

Stose, A. J. and Stose, G. W., 1957, Geology and mineral resources of the Gossan lead district and adjacent areas: Virginia Division of Mineral Resources Bulletin 72, 233 p.

Swain, L. A., Hollyday, E. F., Daniel, C. C., III, Zapecza, O. S., Mesko, T. O., and Wright, W. G., 1991, Plan of study for the regional aquifer-system analysis of the Appalachian Valley and Ridge, Piedmont, and Blue Ridge physiographic provinces of the eastern and southeastern United States, with a description of study-area geology and geohydrology: U.S. Geological Survey Water-Resources Investigations Report 91-4066, 54 p.

Swartz, F. M., 1948, Trenton and sub-Trenton of outcrop areas in New York, Pennsylvania, and Maryland: American Association of Petroleum Geologists Bulletin, v. 32, p. 1493-1595.

Taber, S., 1943, Perennially frozen ground in Alaska: its origin and history: Geological Society of America Bulletin, v. 54, p. 1433-1548.

Tarr, W. S., 1898, The peneplain: American Geologist, v. 21, p. 351-370.

Taylor, F. B., Dion, J. A., and Collins, J. J., 1986, Drinking water quality and acid rain in the eastern United States: Water Quality Bulletin, v. 11, p. 50-57.

Thomas, W. A., 1977, Evolution of the Appalachian-Ouachita salients and recesses from reentrants and promontories in the continental margin: American Journal of Science, v. 277, p. 1233-1278.

Thornbury, W. D., 1954, Principles of Geomorphology: New York, John Wiley and Sons, 618 p.

Thornbury, W. D., 1965, Regional geomorphology of the United States: New York, John Wiley and Sons, 609 p.

Thornbury, W. D., 1969, Principles of Geomorphology, 2nd ed.: New York, John Wiley and Sons, 594 p.

Tiffney, B. H., 1985, The Eocene North Atlantic land bridge: its importance in Tertiary and modern phytogeography of the northern hemisphere: Journal of the Arnold Arboretum, v. 66, p. 243-273.

Troll, C., 1948, Der subnivale oder periglaziale Zyklus der Denudation: Erdkunde, v. 2: p. 1-21.

Tschudy, R. H., 1965, An Upper Cretaceous deposit in the Appalachian Mountains: U.S. Geological Survey Professional Paper 525-B: p. B64-B68.

U.S. Geological Survey, 1991, Water resources data for Pennsylvania, water year 1990--volume 2; U.S. Geological Survey Water-Data Report PA-90-2, 274 p.

Walcott, C. D., 1891, Correlation papers, Cambrian: U.S. Geological Survey Bulletin 81, 447 p.

Walcott, C. D., 1892a, Notes on the Cambrian rocks of Virginia and the southern Appalachians: American Journal of Science, v. 44, p. 52-57.

Walcott, C. D., 1892b, Notes on the Cambrian rocks of Pennsylvania and Maryland from the Susquehanna to the Potomac: American Journal of Science, v. 44, p. 669-682.

Walcott, C. D., 1893, The geologist at Blue Mountain, Maryland: National Geographic Magazine, v. 5, p. 84-88.

Walcott, C. D., 1894a, Notes on the Cambrian rocks of Pennsylvania from the Susquehanna to the Delaware: American Journal of Science, v. 47, p. 309-311.

Walcott, C. D., 1894b, On the Occurrence of <u>Olenellus</u> in the Green Pond Mountain series of northern New Jersey, with a note on the conglomerates: American Journal of Science, v. 47, p. 309-311.

Walcott, C. D., 1894c, Paleozoic intraformational conglomerates: U.S. Geological Survey Bulletin 134, p. 1-43.

Walcott, C. D., 1896, The Cambrian rocks of Pennsylvania: U.S. Geological Survey Bulletin 134, 43 p. Walker, J. E., 1966, Hopewell village, a social and economic history of an iron-making community: Philadelphia, University of Pennsylvania Press, 526 p.

Wallace, P. A. W., 1961, Indians in Pennsylvania: The Pennsylvania Historical and Museum Commission, 138 p. Washburn, A. L., 1980, Geocryology: New York, John Wiley and Sons, 406 p.

Washburn, A. L., 1985, Periglacial problems, in Church, M. and Slaymaker, O., eds., Field and Theory.

Lectures in Geocryology: Vancouver, University of British Columbia Press, p. 166-202.

Watts, W. A., 1979, Late Quaternary vegetation of central Appalachia and the New Jersey Coastal Plain: Ecological Monographs, v. 49, p. 427-469.

Way, J. H., 1986, Your guide to the geology of the Kings Gap area, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Environmental Geology Report 8, 31 p.

Webb, J. R., Cosby, B. J., Galloway, J. N., Hornberger, G.M., and Ryan, P. F., 1989a, The Shenandoah watershed study: An overview: Department of Environmental Sciences, University of Virginia, Charlottesville, VA.

Webb, J. R., Cosby, B. J., Galloway, J. N., Hornberger, G.M., and Ryan, P. F., 1989b, Acidification of native brook trout streams in Virginia, a report to the Virginia Department of Game and Inland Fisheries: The Virginia Trout Stream Sensitivity Study, Project Number F-72.

Whisonant, J. C., 1970, Paleocurrents in basal Cambrian rocks of eastern Tennessee: Geological Society of America Bulletin, v. 81, p. 2781-2786.

Whitaker, J. C., 1955a, Direction of current flow in some Lower Cambrian clastics in Maryland: Geological Society of America Bulletin, v. 66, p. 763-766.

Whitaker, J.C., 1955b, Geology of Catoctin Mountain, Maryland and Virginia: Geological Society of America Bulletin, v. 66, p. 435-462.

White, S. E., 1976, Rock glaciers and block fields, review and new data: Quaternary Research, v. 6, p. 77-97.

Wilden, R. and Mabey, D. R., 1961, Giant desiccation fissures on the Black Rock and Smoke Creek Deserts, Nevada: Science, v. 133, p. 1359-1360.

Wilderman, C.C., 1990, Patterns of variation of pH and alkalinity in Pennsylvania streams, based on data obtained by the Alliance for Acid Rain Monitoring (ALLARM), a citizen's monitoring program, in Proceedings of the Conference on Atmospheric Deposition in Pennsylvania: A Critical Assessment, September 11-14, 1989: Environmental Resources Research Institute, The Pennsylvania State University, University Park, PA., p. 94.

Wilderman, C. C., Cole, C. A., and Nguyen, Q. T., in press, Quality trends of some public water supplies in Pennsylvania: Northeastern Geology.

Wilderman, C. C. and Reuss, C. L., 1991, Analysis of patterns and causes of variation in pH and alkalinity concentrations in Pennsylvania streams, using data collected by the Alliance for Acid Rain Monitoring (ALLARM): Pennsylvania Academy of Science Journal, v. 64, p. 210.

Williams, P. J. and Smith, M. W., 1989, The Frozen Earth: Fundamentals of Geocryology: Cambridge, Cambridge University Press, Studies in Polar Research, 306 p.

Wilshusen, J. P. and Sevon, W. D., 1981, Giant Ripples at Mount Cydonia: Pennsylvania Geology, v. 12, p. 2-8.

Wilshusen, J. P. and Sevon, W. D., 1982, Mount Cydonia Sand Company pit, <u>in</u> Potter, N., Jr., ed., South Mountain: Harrisburg Area Geological Society, Guidebook, p. 28-32.

Wolfe, J. A., 1985, Distribution of major vegetational types during the Tertiary, <u>in</u> Sundquist, E. T. and Broecker, W. S., eds., The carbon cycle and atmospheric CO₂: natural variation Archean to present: American Geophysical Union Monograph 32, p. 357-375.

Wood, G. D. and Clendening, J. A., 1982, Acritarchs from the Lower Cambrian Murray Shale, Chilhowee Group of Tennessee, U.S.A.: Palynology, v. 6, p. 255-265.

Woodward, N. B., 1985, Valley and Ridge thrust belt: balanced structural sections Pennsylvania to Alabama: University of Tennessee, Department of Geological Sciences, Studies in Geology 12, 64 p.

Worsley, P., 1984, Periglacial environment: Progress in Physical Geography, v. 8, p. 270-276.

Zietz, I., Haworth, R. T., Williams, H., and Daniels, D. L., 1980, Magnetic anomaly map of the

Appalachians: Memorial University of Newfoundland, Map No. 2, scale 1:1,000,000.