


Guidebook

# *56th Annual Field Conference of Pennsylvania Geologists*

## Geology in the South Mountain Area, Pennsylvania



Hosts:  
Dickinson College  
Pennsylvania Geological Survey

September 26, 27, and 28, 1991  
Carlisle, Pa.

Guidebook for the  
56th ANNUAL FIELD CONFERENCE OF PENNSYLVANIA GEOLOGISTS  
**GEOLOGY IN THE SOUTH MOUNTAIN AREA, PENNSYLVANIA**

W. D. Sevon and Noel Potter, Jr., editors

Contributors and field trip leaders:

Rolf, Ackermann  
John H. Barnes  
Albert E. Becher  
S. W. Berkheiser, Jr.  
Douglas C. Chichester  
G. Michael Clark  
Gretchen Dockter  
Henry W. A. Hanson  
Marcus M. Key, Jr.  
Steven Lev  
Laura Pezzoli  
Robert J. Schott  
Samuel J. Sims  
Samuel I. Root  
R. C. Smith, II  
Thomas Troy  
Randal L. Van Scyoc  
John H. Way  
Candie C. Wilderman  
Ellis L. Yochelson

September 26-28, 1991

Hosts: Dickinson College  
Pennsylvania Geological Survey

Headquarters Motel: The Embers Convention Center, Carlisle, PA

Cover: Topographic maps at 1 inch = 1,600 feet for 3 stops on the field trip. Older maps are from a series with a contour interval of 10 feet that cover the entire South Mountain area in Pennsylvania. They were surveyed and drafted by Ambrose E. Lehman 1875-1885, and appear in Atlases D5 and D6 of the Second Pennsylvania Geological Survey. Modern maps are:

Stop 3, USGS Iron Springs 7.5-minute quadrangle, C.I. = 20 feet.  
Stop 6, USGS Scotland and Caledonia Park 7.5-minute quadrangle, C.I. = 20 feet.

Stop 8, USGS Dickinson and Mount Holly Springs 7.5-minute quadrangle, C.I. = 10 feet.

Guidebooks distributed by:

Field Conference of Pennsylvania Geologists, Inc.  
P. O. Box 1124  
Harrisburg, PA 17108

# THE LOWER CAMBRIAN CLASTICS OF SOUTH MOUNTAIN, PENNSYLVANIA

by

Marcus M. Key, Jr.

Department of Geology, Dickinson College, Carlisle, PA

## INTRODUCTION

The northeast/southwest trending linear ridges of South Mountain are composed of the Lower Cambrian clastic metasedimentary rocks of the Chilhowee Group (Figure 3). The group comprises four formations (from oldest to youngest): Loudoun, Weverton, Harpers, and Antietam. In the South Mountain area, the thickness of the group ranges from roughly 3,000 feet in the Mt. Holly area to 5,000 feet in the Caledonia area. The group consists of lithogenic sediments which originally were shales, litharenites, sublitharenites, quartz arenites, and conglomerates that have been metamorphosed into phyllites, quartzites, and squashed-pebble conglomerates. Currently only the Montalto Member of the Harpers Formation and the Antietam Quartzite are economically important as a source of coarse and fine aggregate, and colored sands for speciality uses.

## AGE

The age of the Chilhowee Group has historically been a troublesome issue (Nickelsen, 1956). The problem existed because of a paucity of datable materials in these rocks. The age of the group is now rather certain to be Early Cambrian based on the radiometrically dated, stratigraphically older Catoctin Formation and two biostratigraphically dated formations in the upper and lower units of the Chilhowee Group.

Until recently, the age of the Catoctin Formation was widely disputed with radiometric dates ranging from 420+4 Ma (Silurian) (Nagel and Mose, 1984) to 820 Ma (Precambrian) (Rankin and others, 1969). Recent radiometric ages have been much more consistent. The metarhyolites in the Catoctin Formation from South Mountain have been dated at 597+18 Ma (Aleinikoff and others, 1991). Badger and Sinha (1988) determined the age of the Catoctin in west-central Virginia to be 570+36 Ma. The stratigraphically older Lynchburg Formation in Virginia has been dated at 600 Ma (Mose and others, 1985). All of these recent dates suggest the Catoctin Formation is latest Precambrian in age.

Biostratigraphic constraints on the age of the Chilhowee Group come from fossils in the upper and lower parts of the group. The Early Cambrian age fossils first reported by Walcott (1891) from the Antietam Quartzite have been found throughout the Appalachians in this formation and its stratigraphic equivalents to the south (See Yochelson, this guidebook). The age of the upper Chilhowee rocks has been repeatedly confirmed with different index fossils and in different areas along the Appalachians. The Antietam Quartzite and its stratigraphic equivalents contain a diverse fauna indicative of the Early Cambrian. This fauna contains acritarchs, the worm tube ichnofossil *Skolithos linear-*




Figure 3. Geologic map of South Mountain area showing distribution of Chilhowee Group formations (Modified from Berg and others, 1980). Numbers refer to Chilhowee Group field trip stops.

is, the inarticulate brachiopod *Obolella*, the articulate brachiopod *Camarella minor*, the trilobite *Olenellus*, the primitive mollusc *Hyolithus communis*, and the ostracod *Indiana tennesseensis*. This fauna has been reported from Pennsylvania to Tennessee (Walcott, 1896; Stose, 1909; Bassler, 1919; Barrell, 1925; Stose and Bascom, 1929; Stose, 1932; Resser and Howell, 1938; Amsden, 1951; Stose, 1953; Stose and Stose, 1957; Laurence and Palmer, 1963; Wood and Clendening, 1982). The occurrence of *Olenellus* constrains the Antietam Quartzite to the *Bonnina-Olenellus* Biozone which existed from 550-540 Ma (Fritz, 1972; Palmer, 1981, 1983). This dates the Antietam Quartzite as latest Early Cambrian.

The presence of *Skolithos linearis* in the Montalto Member of the Harpers Formation does not provide any biostratigraphic control because this trace fossil occurs in both Cambrian and Precambrian deposits (Fritz and Crimes, 1985). Recently an age diagnostic fauna has been found in the stratigraphic equivalent of the Weverton Quartzite in southwestern Virginia (Simpson and Sundberg, 1987). This depauperate fauna contains many small shelly fossils and the trace fossil *Rusophycus* which is indicative of earliest Early Cambrian (Fedonkin, 1981). This constrains the age of the lower Chilhowee Group to 570-560 Ma (Sepkoski and Knoll, 1983).

The radiometric dates of the Catoctin Formation place a maximum age on the Chilhowee Group of roughly 570 Ma. The biostratigraphic data suggest an age range of 570-540 Ma. Based on the stratigraphic conformity of all of the Chilhowee Group formations (Nickelsen, 1956; Fauth, 1968), the age of the entire group is considered to be Early Cambrian.

#### CATOCTIN FORMATION - CHILHOWEE GROUP CONTACT

Underlying the Chilhowee Group is the Catoctin Formation (See the paper by Smith and others, this guidebook). The Catoctin consists of aphanitic, extrusive volcanics (dominantly basalts and rhyolites) that have been subsequently metamorphosed. These represent late Precambrian rift-related volcanism (Badger and Sinha, 1991) which occurred during the opening of the Proto-Atlantic (Iapetus) Ocean (Rankin, 1975). They are inferred to have been generally extruded subaerially based on the absence of pillow lavas and the presence of columnar jointing (Reed, 1955; Fauth, 1968). Petrographic, rare earth element, and major element analyses have confirmed this (Rankin and others, 1969; Dockter, 1990). However, some local subaqueous extrusion has been documented by the presence of pillow lavas in Virginia (Bowring and Spencer, 1987; Kline and others, 1987).

The Catoctin-Chilhowee Group (i.e., Catoctin-Loudoun) contact is not exposed in the South Mountain area (Stose, 1909; Stose and Bascom, 1929; Fauth, 1968). Historically, the contact has been considered both conformable (Bloomer and Bloomer, 1947; Bloomer, 1950; Cloos, 1951; Rankin, 1967) and unconformable (Stose, 1906, 1909, 1932; Stose and Bascom, 1929; King, 1950; Stose and Stose, 1957; King and Fergusen, 1960; Rankin and others, 1969). The absence of a structural discordance between the Catoctin and Loudoun Formations suggests the contact is conform-

able (Bloomer, 1950; Bloomer and Werner, 1955; Reed, 1955; Fauth, 1968). Both formations exhibit similar deformation patterns (i.e., orientation of cleavage and lineations) (Stose, 1932; King, 1949; Cloos, 1951; Freedman, 1967; Fauth 1968). If there was a period of deformation prior to the deposition of the Loudoun Formation, it would indicate unconformity. Bloomer (1950), Cloos (1951), and Bowring and Spencer (1987) argue that the contact is also conformable due to the interfingering of the Catoctin volcanics with the Loudoun clastics in Virginia.

The presence of Catoctin fragments in the basal clastic sediments of the Loudoun Formation suggests the contact is unconformable [But see Bloomer (1950) for an opposing view]. This indicates the Catoctin was subaerially exposed and eroded prior to deposition of the Loudoun (Freedman, 1967). According to Reed (1955) this period of erosion is preserved by a metamorphosed saprolite at the contact. Another argument for unconformity is that the contact represents a change from terrestrial extrusion of the Catoctin to submarine deposition of the Loudoun (Nickelsen, 1956; Fauth, 1968).

The contact may be unconformable, but it probably represents only a brief time hiatus (Badger and Sinha, 1988). The previously discussed radiometric ages of the Catoctin and the biostratigraphic ages of the Chilhowee suggest little if any time is missing at this contact.

#### LITHOSTRATIGRAPHY

The basal Chilhowee unit is the Loudoun Formation. The sediments of this formation contain fragments of the underlying Catoctin metarhyolites (Stose, 1906, 1909, 1932; Stose and Bascom, 1929; Freedman, 1967; Fauth, 1968). Thickness in the South Mountain area ranges from 0 to 550 feet (Stose and Bascom 1929, Stose 1932, Freedman 1967, Fauth 1968). In the Caledonia and Mt. Holly areas, this unit crops out along the southeast slope of Piney Mountain. The lithology ranges from a purplish phyllite to a grayish, quartz/phyllite/rhyolite-pebble metaconglomerate (Stop 9). The grains are subrounded to subangular grains with fair sorting (Fauth, 1968).

Overlying the Loudoun Formation is the Weverton Quartzite whose thickness in the South Mountain area ranges from 600 to 1,400 feet (Stose and Bascom, 1929; Stose, 1932; Freedman, 1967; Fauth 1968). In the Caledonia area, this unit crops out along the crests of Piney Mountain and Big Pine Flat Ridge and along the southeast slope of East Big Flat Ridge. In the Mt. Holly area, it crops out along the crest of Mt. Holly (including Hammond's Rocks, Stop 8) and along the southeast slope of Piney Mountain. The lithology ranges from grayish-green and purplish quartzites and quartzose graywackes with occasional thin zones of graywacke conglomerate, quartzite, and phyllite. The grains are subrounded to subangular with moderate sorting (Fauth, 1968).

Overlying the Weverton Quartzite is the Harpers Formation whose thickness in the South Mountain area ranges from 1,500 to 3,000 feet (Stose and Bascom, 1929; Stose, 1932; Freedman, 1967; Fauth, 1968). In the Caledonia area, this unit crops out along

the crests of Piney Mountain and Big Pine Flat Ridge. In the Mt. Holly area, it crops out along the crest of Piney Mountain and around the nose of Mt. Holly. There are two distinct lithologies present. The lower light-gray quartzite has good sorting with subrounded grains (Fauth, 1968). This unit is called the Montalto Quartzite Member, and it is 1,200 to 2,500 feet thick. The Montalto is crossbedded and has numerous *Skolithos linearis* worm tubes (Figure 2). In the upper part of this member is a distinctive bluish quartzite unit that ranges from 15 to 25 feet thick and is useful in mapping (Fauth, 1968). The upper greenish gray-wacke/phyllite is moderately sorted with subrounded grains. It is 300 to 500 feet thick (Fauth, 1968; Root, 1978).



Figure 2. *Skolithos linearis* worm tubes from the Antietam Formation.

Overlying the Harpers Formation is the youngest Chilhowee unit, the Antietam Quartzite whose thickness in the South Mountain area ranges from 440 to 900 feet (Stose, 1909; Stose and Bascom, 1929; Stose, 1932; Freedman, 1967; Fauth, 1968; Root, 1978). The Pennsylvania Geological Survey is currently studying core from an angle hole that penetrated the entire thickness of the Antietam north of Caledonia Park. In the Caledonia area, this unit crops out around the nose of Big Pine Flat Ridge. In the Mt. Holly area, it crops out along the northwest slope and the nose of Mt. Holly. The lithology is dominated by white to grayish quartzite which is medium- to coarse-grained, subrounded to rounded, with good sorting (Fauth, 1968). *Skolithos linearis* worm tubes (Figure 2) are abundant with some up to 15 inches long (Shirk, 1980; Wilshusen and Sevon, 1981, 1982). Megaripples are present in the Mt. Cydonia area (Wilshusen and Sevon, 1981, 1982) (Stop 5). There are numerous abandoned aggregate pits and quarries developed in the Antietam in the South Mountain area. Significant active aggregate operations in the Antietam occur at Mt. Cydonia (Stop 5) and Mt. Holly Springs (Stop 11) as well as Toland.

## CHILHOWEE GROUP - TOMSTOWN DOLOMITE CONTACT

Overlying the Chilhowee Group is a thick sequence of Cambrian through Ordovician carbonates. The oldest of these carbonates is the Tomstown Dolomite which rests upon the Antietam Quartzite. The Tomstown is dominantly dolomitic with some minor limestone. The contact between the Antietam Quartzite and the Tomstown Dolomite is apparently gradational as evidenced by a calcareous basal shale in the Tomstown (Brezinski, 1991). In the South Mountain region, this contact is marked by the presence of kaolinite-rich white clays (Berkheiser and others, 1982). This clay formed either from dissolution of the Tomstown which resulted in the concentration of its contained lithogenic sediment (Stose, 1907) or from hydrothermal alteration of the Tomstown (Hosterman, 1968, 1969).

The contact is often assumed to be conformable even though it is not exposed in the South Mountain area (Stose, 1909; Stose and Bascom, 1929; Fauth, 1968; Freedman, 1968; Brezinski, 1991). Besides the lithologic argument for conformity mentioned above, there is also biostratigraphic support for conformity. The Tomstown Dolomite has been dated as latest Early Cambrian due to the presence of the trilobite *Olenellus* (Fauth, 1968). As the Tomstown and Antietam are of the same age based on the available biostratigraphic control, the contact is probably conformable (Fauth, 1968).

## PALEOGEOGRAPHY

The Chilhowee Group was deposited in rift-induced sedimentary basins that formed as a result of the opening of the Iapetus Ocean in the late Precambrian (Bond and others, 1984). In the Early Cambrian, clastic sediments (including the Chilhowee Group) were deposited along the newly-formed southern and eastern edges of the North American craton (Thomas, 1977). These sediments represent the basal transgressive sediments of the Sauk Sequence (Sloss, 1963; Brown, 1970; Schwab, 1972). The sediments were eroded from the craton, transported to the southeast (Kay, 1951; Whitaker, 1955; Brown, 1970; Schwab, 1970; Whisonant, 1970), and were deposited in shallow marine environments of the Iapetus Ocean along the coast of North America. During this time, the South Mountain area was located at roughly 15° south latitude (Scotese and others, 1979).

Deposition of the Chilhowee Group began with the transgression of the Cambrian sea which onlapped from the southeast and inundated the subaerially exposed Precambrian Catoctin Formation (Nickelsen, 1956; Fauth, 1968). The first sediments resulted in the Loudoun Formation which was partially derived from and deposited on the Catoctin Formation. This is evidenced by the clasts of Catoctin metarhyolites which were incorporated into the basal Loudoun sediments.

Evidence of a northwestern cratonic source area for the Chilhowee Group comes from stratigraphic thicknesses, paleocurrent indicators, and facies distributions. The Loudoun Formation and the Weverton Quartzite thicken to the west (Swartz,

1948). The Montalto Quartzite Member of the Harpers Formation thickens to the north (Fauth, 1968). Stose and Bascom (1929) reported that the entire Chilhowee Group thickens to the northwest. Crossbedding in the Weverton Quartzite indicates a western source area (Whitaker, 1955), while ripple marks in the Antietam Quartzite suggest a northwestern source area (Wilshusen and Sevon, 1981, 1982). The upper Chilhowee formation grades eastwardly from the Antietam Quartzite in the Blue Ridge Mountains to the Chickies quartz-mica schist in the Piedmont (Freedman, 1968). This facies change suggests deeper water to the east and a westward source area.

The depositional environments represented by the Chilhowee Group in the South Mountain area are dominantly shallow marine on continental shelf environments (Nickelsen, 1956). This is evidenced by the presence of trilobites, brachiopods, ostracods, and worm tubes. Water depth increased to the southeast toward the deep-marine Wissahicken basin (Thomas, 1977) where the Chilhowee Group was deposited below storm-wave base (Freedman, 1968; Simpson and others, 1991).

The depositional environment of the Antietam Quartzite can be further refined as a shallow marine beach setting. The Antietam's clean, well-rounded quartz sand and the presence of mega-ripples led Wilshusen and Sevon (1981, 1982) to interpret this formation as reflecting a near-shore to offshore, below normal wave-base setting. Based on the outcrop pattern and the morphology of the sand bodies, Kauffman and Frey (1979) inferred that the Antietam was deposited in a barrier island environment. The occurrence of *Skolithos linearis* worm tubes in the Antietam places this formation in the *Skolithos* ichnofacies. This facies is indicative of the intertidal to subtidal zone of a shallow, high energy, coastal, beach environment (Seilacher, 1967; Crimes, 1970). The Montalto Quartzite Member of the Harper Formation has identical worm tubes, and it is interpreted to have a similar depositional environment.

Fluctuations in local relief, rate of uplift, and rate of erosion in the source area relative to the rate of subsidence and deposition in the depositional basins resulted in local fining and coarsening trends within the Chilhowee Group. On a regional scale over the course of Chilhowee deposition, sediments generally became finer as well as texturally and mineralogically more mature in response to rising sea level and increasing tectonic stability of the east coast of the North American craton (Fauth, 1968; Patterson and Simpson, 1991). The maturing upward trend is evidenced by a general increase in the degree of sorting and roundness as well as an increase in the amount of quartz clasts relative to those of feldspar (Fauth, 1968). In the South Mountain area, this can be grossly seen by comparing the immature Loudoun conglomerates (Stop 9) with the mature Antietam quartz arenites (Stops 5 and 11). The fining upward trend continued after Chilhowee deposition into the Tomstown Dolomite. Before the end of the Early Cambrian, the sediment became finer until little or no lithogenic sediment was entering the sea and carbonate biogenic sedimentation was initiated (Freedman, 1967, 1968).

# REFERENCES CITED

- Adams, G. F., ed., 1975, Planation surfaces. peneplains, pediplains, and etchplains: Harrisburg, Dowden, Hutchinson and Ross, Inc., 476 p.
- Aleinikoff, J. N., Zartman, R. E., Rankin, D. W., Lyttle, P. T., Burton, W. C., and McDowell, R. C., 1991, New U-Pb zircon ages for rhyolite of the Catoctin and Mount Rogers Formations-More evidence for two pulses of Iapetan rifting in the central and southern Appalachians (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 2.
- Allen, J. R. L., 1982, Sedimentary structures, their character and physical basis: New York, Elsevier Publishing Co., 1256 p.
- Amsden, T. W. 1951, Paleontology of Washington County: Maryland Department of Geology and Mines, Water Resources Bulletin, No. 10, p. 98-123.
- Anonymous, We. R., 1985, Gross Minerals, a force in the fine phyllite fillers field, Pennsylvania Geology, v. 16, no. 6, p. 2-5.
- Badger, R. L. and Sinha, A. K., 1988, Age and Sr isotopic signature of the Catoctin volcanic province: Implications for subcrustal mantle evolution: Geology, v. 16, p. 692-695.
- Badger, R. L. and Sinha, A. K., 1991, Nature of late Precambrian mafic magmatism within the Appalachian orogen (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 5.
- Barrell, J., 1925, The nature and environment of the Lower Cambrian sediment of the southern Appalachians: American Journal of Science, v. 9, p. 1-20.
- Barron, E. J., 1989, Climate variations and the Appalachians from the Late Paleozoic to the Present: results from model simulations, in Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 99-118.
- Bascom, F., 1893, The structures, origin, and nomenclature of the acid volcanic rocks of South Mountain, Pennsylvania: Journal of Geology, v. 1, p. 813-832.
- Bassler, R. S., 1919, Cambrian and Ordovician: Maryland Geological Survey, Johns Hopkins University Press, Baltimore, 408 p.
- Baulig, H., 1952, Surfaces d'Aplanissement: Annales de Géographie, v. 61, p. 161-183; 245-262.
- Becher, A. E. and Root, S. I., 1981, Groundwater and geology of the Cumberland Valley, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Water Resource Report 50, 95 p.
- Becher, A. E. and Taylor, L. E., 1982, Groundwater resources in the Cumberland and contiguous valleys of Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Water Resource Report W-53, 67 p.
- Berg, T. M., 1975, Geology and mineral resources of the Brodheadsville quadrangle, Monroe and Carbon Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 205a, 60 p.
- Berg, T. M., and others, 1980, Geologic Map of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Map 1, scale 1:250,000.
- Berg, T. M., Sevon, W. D., and Abel, R., 1984, Rock types of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Map 63, scale 1:500,000.
- Berkheiser, S. W., Jr., 1985, High-purity silica occurrences in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 88, 67 p.
- Berkheiser, S. W., Jr., Barnes, J. H., and Smith, R. C., II, 1985, Directory of the nonfuel-mineral producers in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Information Circular 54, 4th ed., 165 p.
- Berkheiser, S. W., Jr., Potter, N., Jr., and Sevon, W. D., 1982, Hempt Brothers clay pit at Toland: in Potter, N., Jr., ed., South Mountain: Guidebook, Harrisburg Area Geological Society, p. 10-14.
- Berkheiser, S. W., Jr. and Smith, R. C., II, 1990, Overhead protection, Pennsylvania's mineral industry contribution: Pennsylvania Geology, v. 21, no. 3, p. 2-6.
- Bining, A. C., 1979, Pennsylvania iron manufacture in the eighteenth century, 2nd ed.: Harrisburg, Pennsylvania Historical and Museum Commission, 215 p.
- Bjornerud, M. G., 1991, Conveying principles of finite strain with standard graphics software: Journal of Geological Education, v. 39, p. 23-27.
- Black, R. F., 1976a, Periglacial features indicative of permafrost: ice and soil wedges: Quaternary Research, v. 6, p. 3-26.
- Black, R. F., 1976b, Features indicative of permafrost: Annual Review of Earth and Planetary Sciences, v. 4, p. 75-94.
- Black, R. F., 1983, Pseudo-ice-wedge casts of Connecticut, northeastern United States: Quaternary Research, v. 20, p. 74-89.
- Blatt, H., 1980, Origin of sedimentary rocks, 2nd ed.: Englewood Cliffs, NJ, Prentice Hall, 782 p.
- Bloomer, R. O., 1950, Late Pre-Cambrian and Lower Cambrian formations in central Virginia: American Journal of Science, v. 248, p. 753-783.
- Bloomer, R. O. and Bloomer, R. R., 1947, The Catoctin Formation in central Virginia: Journal of Geology, v. 55, p. 94-106.
- Bloomer, R. O. and Werner, H. J., 1955, Geology of the Blue Ridge in central Virginia: Geological Society of America Bulletin, v. 65, p. 579-606.

- Bond, G. C., Nickeson, P. A., and Kominz, M. A., 1984, Breakup of a supercontinent between 625 Ma and 555 Ma: New evidence and implications for continental histories: *Earth and Planetary Science Letters*, v. 70, p. 325-345.
- Bowring, C. and Spencer, E., 1987, Catoctin pillow lavas (abs.): *Geological Society of America Abstracts with Programs*, v. 19, no. 2, p. 77.
- Bradt, P. T. and Wimpfheimer, L., 1987, Survey of Pocono lakes to evaluate sensitivity to acidification: *Environmental Studies Center, Lehigh University, Bethlehem, PA.*
- Braun, D. D., 1989a, A revised Pleistocene glaciation sequence in eastern Pennsylvania: Support for limited Early Wisconsin ice and a single Late Illinoian advance beyond the Late Wisconsin border (abs.): *28th International Geological Congress Abstracts*, v. 1: p. 1-196-1-197.
- Braun, D. D., 1989b, Glacial and periglacial erosion of the Appalachians, in Gardner, T. W. and Sevon, W. D., eds., *Appalachian Geomorphology* (reprinted from *Geomorphology*, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 233-256.
- Braun, D. D., and Inners, J. D., 1990, Weathering of conglomerate ledges and tors within the glacial limit in northeastern Pennsylvania: Evidence for single stage tor development under Pleistocene periglacial conditions (abs.): *Geological Society of America Abstracts with Programs*, v. 22, no. 2, p. 6.
- Braun, E. L., 1950, *Deciduous Forests of Eastern North America*: Philadelphia, Blackiston Company, 596 p.
- Brezinski, D. K., 1991, Lithostratigraphy of the Lower Cambrian Tomstown Formation (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 1, p. 10.
- Brown, W. R., 1970, Investigations of the sedimentary record in the Piedmont and Blue Ridge of Virginia, in Fischer, G. W., Pettijohn, F. J., Reed, J. C., Jr., and Weaver, K. N., eds., *Studies of Appalachian Geology: Central and Southern*: New York, Interscience, p. 335-349.
- Bryan, K., Cleaves, A. B., and Smith, H. T. U., 1932/33, The present status of the Appalachian problem: *Zeitschrift für Geomorphologie*, v. 7, p. 312-320.
- Bryan, K., 1946, Cryopedology the study of frozen ground and intensive frost-action with suggestions on nomenclature: *American Journal of Science*, v. 244, p. 622-642.
- Büdel, J., 1982, *Climatic Geomorphology*: Princeton, Princeton University Press, 443 p.
- Camuto, C., 1991, Dropping acid in the Southern Appalachians: A wild trout resource at considerable risk: *Trout, Winter*, 1991, p. 16-39.
- Carter, B. J., and Ciolkosz, E. J., 1980, Soil temperature regimes of the Central Appalachians: *Soil Science Society of America Journal*, v. 44, p. 1052-1058.
- Carter, B. J., and Ciolkosz, E. J., 1986, Sorting and thickness of waste mantle material in a sandstone spur in central Pennsylvania: *Catena*, v. 13, p. 241-256.
- Cecil, C. B., 1990, Paleoclimatic controls on stratigraphic repetition of chemical and siliclastic rocks: *Geology*, v. 19, p. 533-536.
- Ciolkosz, E. J., Carter, B. J., Hoover, M. T., Counce, R. C., Waltman, W. J., and Dobos, R. R., 1990, Genesis of soils and landscapes in the Ridge and Valley province of central Pennsylvania, in Knuepfer, P. L. K. and McFadden, L. D., eds., *Soils and landscape evolution*: Amsterdam, Elsevier, p. 245-261.
- Ciolkosz, E. J., Counce, R. C., and Sevon, W. D., 1986, *Periglacial features in Pennsylvania*: University Park, PA, The Pennsylvania State University, Agronomy Department, Agronomy Series Number 92, 15 p.
- Clark, G. M., 1989a, Central and southern Appalachian accordant ridge crest elevations south of glacial border: Regional erosional remnants or incipient periglacial surfaces? (abs.): *28th International Geological Congress Abstracts*, v. 1, p. 1-299-1-300.
- Clark, G. M., 1989b, Central and southern Appalachian water and wind gap origins: Review and new data, in Gardner T. W. and Sevon, W. D., eds., *Appalachian Geomorphology* (reprinted from *Geomorphology*, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 209-232.
- Clark, G. M., and Ciolkosz, E. J., 1988, Periglacial geomorphology of the Appalachian Highlands and Interior Highlands south of the glacial border: *Geomorphology*, v. 1, p. 191-220.
- Clark, G. M., Ciolkosz, E. J., Kite, J. S., and Lietzke, D. A., 1989, *Central and Southern Appalachian Geomorphology - Tennessee, Virginia, and West Virginia*: 28th International Geological Congress Field Trip Guidebook T150, 105 p.
- Clark, G. M., and Hedges, J., in press, Origin of certain high-elevation local broad uplands in the Central Appalachians south of the glacial border - a paleoperiglacial hypothesis: *21st Annual Binghamton Geomorphology Symposium*, Chichester, Wiley, in press.
- Clark, M. J., ed., 1988, *Advances in periglacial geomorphology*: New York, John Wiley and Sons, 481 p.
- Clarke, M. S., 1968, *Pioneer iron works*: Philadelphia, Chilton Book Company, 80 p.
- Cleaves, E. T., Edwards, J. E., Jr., and Glaser, J. D., 1968, *Geologic map of Maryland*: Maryland Geological Survey, scale 1:250,000.
- Cloos, E., 1947, Oolite deformation in the South Mountain fold, Maryland: *Geological Society of America Bulletin*, v. 58, p. 843-918.
- Cloos, E., 1951, *Physical features of Washington County*: Maryland Department of Geology and Mines, *Water Resources Bulletin* 10, p. 17-94.
- Cloos, E., 1958, *Structural geology of South Mountain and Appalachians in Maryland*: 23rd Annual Field Conference of Pennsylvania Geologists, Hagerstown, MD, Guidebook, 85 p.

- Cloos, E. and others, 1968, The geology of mineral deposits in south-central Pennsylvania: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, 80 p.
- Cloos, E., 1971, Microtectonics along the western edge of the Blue Ridge, Maryland and Virginia: Johns Hopkins University, Studies in Geology No. 20, 234 p.
- Cosby, B. J., Hornberger, G. M., Wright, R. F., and Galloway, J. N., 1986, Modeling the effects of acid deposition: control of long-term sulfate dynamics by soil sulfate adsorption: *Water Resources Research*, v. 22, p. 1283-1291.
- Crimes, T. P., 1970, The significance of trace fossils in sedimentology, stratigraphy and palaeoecology with examples from Lower Palaeozoic strata, *in* Crimes, T. P. and Harper, J. C., eds., *Trace Fossils: Geological Journal Special Issue 3*, p. 106-126.
- Daly, R. A., 1905, The accordance of summit levels among Alpine mountains: the fact and its significance: *Journal of Geology*, v. 13, p. 105-125.
- Davis, W. M., 1889, The rivers and valleys of Pennsylvania: *National Geographic Magazine*, v. 1, p. 183-253.
- Delcourt, P. A. and Delcourt, H. R., 1981, Vegetation maps for eastern North America; 40,000 yr BP to the present, *in* Romans, R. C., ed., *Geobotany II: New York*, Plenum, p. 123-165.
- Delcourt, P. A. and Delcourt, H. R., 1984, Late Quaternary paleoclimates and biotic responses in eastern North America and the western North Atlantic Ocean: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 48: p. 263-284.
- Demek, J., 1969, Cryoplanation terraces, their geographical distribution, genesis and development: *Rozprawy Československé Akademie Věd. Rada Matematických a Přírodních Věd*, v. 79, no. 4, 80 p., 16 photos.
- Demek, J., 1972, *Manual of Detailed Geomorphological Mapping: Prague*, Academia, 344 p.
- d'Inwilliers, E. V., 1887, Report on the iron ore mines and limestone quarries of Cumberland-Lebanon Valley, *in* Annual report of the Geological Survey for 1886: Pennsylvania Geological Survey, 2nd ser., Annual Report 1886, Pt. IV, p. 1411-1567.
- Dockter, G. N., 1990, The Catoctin metabasalts: evidence of a rifting environment: unpublished independent research paper, Dickinson College, Carlisle, PA, 27 p.
- Duffy, D. F. and Wittecar, G. R., 1991, Geomorphic development of segmented alluvial fans in the Shenandoah Valley, Stuarts Draft, Virginia (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 1, p. 24.
- Edwards, J., Jr., compiler, 1978, *Geologic map of Washington County: Maryland Geological Survey*, scale 1:62,500.
- Ehlen, J., 1990, Geomorphic, petrographic and structural classification of granite landforms using spatial patterns (abs.): *Geological Society of America Abstracts With Programs*, v. 22, no. 7, p. A21.
- Espenshade, G. H., 1986, *Geology of the Marshall quadrangle, Fauquier County, Virginia: U. S. Geological Survey Bulletin 1560*, 59 p.
- Faill, R. T. and MacLachlan, D. B., 1989, Tectonic terranes of southeastern Pennsylvania (abs.): *Geological Society of America Abstracts with Programs*, v. 21, no. 1, p. 13.
- Farlekas, G., 1961, The geology of part of South Mountain of the Blue Ridge province north of the Pennsylvania-Maryland border: unpublished M.S. thesis, University Park, The Pennsylvania State University, 64 p.
- Fauth, J. L., 1968, *Geology of the Caledonia Park quadrangle Area, South Mountain, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 129a*, 133 p.
- Fauth, J. L., 1977, *Geologic map of the Catoctin Furnace and Blue Ridge Summit Quadrangles, Maryland: Maryland Geological Survey*, scale 1:24,000.
- Fauth, J. L., 1978, *Geology and mineral resources of the Iron Springs area, Adams and Franklin Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 129c*, 72 p.
- Fauth, J. L., 1981, *Geologic map of the Myersville Quadrangle, Maryland: Maryland Geological Survey*, scale 1:24,000.
- Fedonkin, M. A., 1981, Palaeoichnology of the Precambrian-Cambrian transition, *in* Taylor, M. E., ed., *Short papers from the Second International Symposium on the Cambrian System: U.S. Geological Survey Open-File Report 81-743*, p. 89-90.
- Fenneman, N. M., 1938, *Physiography of eastern United States: New York*, McGraw-Hill Book Company, 714 p.
- Flemal, R. C., 1971, The attack on the Davisian system of geomorphology: a symposium: *Journal of Geologic Education*, v. 19, p. 3-13.
- Flinn, D., 1962, On folding during three-dimensional progressive deformation: *Geological Society of London Quarterly Journal*, v. 118, p. 385-433.
- Flippo, H. N., Jr., 1974, *Springs of Pennsylvania: Pennsylvania Department of Environmental Resources, Water Resources Bulletin 10*, 46 p.
- Flower, L. E., 1975, *History of Pine Grove Furnace: Carlisle, PA, Cumberland County Historical Society*, 21 p.
- Foose, R. M., 1945a, *Iron-manganese ore deposits of White Rocks, Cumberland County: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 26*, 35 p.
- Foose, R. M., 1945b, *Manganese minerals of Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resource Report 27*, 130 p.
- Foose, R. M. and others, 1948, *Excursion 1, South Mountain, in* Cathcart, S. H., ed., *14th Annual Field*

- Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 1-6.
- Frakes, L. A., 1979, *Climates throughout geologic time*: Amsterdam, Elsevier Science Publishers, 310 p.
- Freedman, J., 1967, *Geology of a portion of the Mount Holly Springs Quadrangle, Adams and Cumberland Counties, Pennsylvania*: Pennsylvania Geological Survey, 4th ser., Progress Report 169, 66 p.
- Freedman, J., 1968, *Bender's Quarry - Mt. Holly Springs, Pennsylvania*, in *The geology of mineral deposits in south-central Pennsylvania*: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 28-37.
- French, H. M., 1976, *The periglacial environment*: London, Longman, 309 p.
- French, H. M. and Karte, J., 1988, *A periglacial overview*, in Clark, M. J., ed., *Advances in Periglacial Geomorphology*: New York, John Wiley and Sons, p. 463-473.
- Friedman, G. M. and Sanders, J. E., 1978, *Principles of Sedimentology*: New York, John Wiley and Sons, 792 p.
- Fritz, W. H., 1972, *Lower Cambrian trilobites from the Sekawi Formation type section, Mackenzie Mountains, north-western Canada*: Geological Survey of Canada Bulletin, v. 212, p. 1-58.
- Fritz, W. H. and Crimes, T. P., 1985, *Lithology, trace fossils, and correlation of the Precambrian-Cambrian boundary beds, Cassiar Mountains, north-central British Columbia*: Geological Survey of Canada Paper 83-13, p. 1-24.
- Fulton, R. J., 1989, ed., *Quaternary Geology of Canada and Greenland*: Geological Society of America, Volume K-1, Geological Survey of Canada, no. 1, 839 p.
- Gagen, C. J. and Sharpe, W. E., 1987, *Net sodium loss and mortality of three salmonid species exposed to a stream acidified by atmospheric deposition*: Environmental Contamination and Toxicology Bulletin, v. 39, p. 7-14.
- Gannett Fleming Corrdry and Carpenter, Inc., 1945, *Report on water treatment project for the Borough of Chambersburg, PA*: Harrisburg, PA.
- Gardiner, V. and Dackombe, R., 1983, *Geomorphological Field Manual*: London, George Allen and Unwin, 254 p.
- Gardner, T. W. and Sevon, W. D., eds., 1989, *Appalachian Geomorphology*: Amsterdam, Elsevier, 318 p.
- Geiser, P. A. and Engelder, T., 1983, *The distribution of layer-parallel shortening fabrics in the Appalachian foreland of New York and Pennsylvania: Evidence for two non-coaxial phases of the Alleghanian orogeny*, in Hatcher, R. D., Jr., Williams, H., and Zeitz, I., eds., *Contributions to the tectonics and geophysics of mountain chains*: Geological Society of America Memoir 158, p. 161-175.
- Gerhart, J. M. and Lazorchick, G. J., 1988, *Evaluation of the ground-water resources of the lower Susquehanna River Basin, Pennsylvania and Maryland*: U.S. Geological Survey Water-Supply Paper 2284, 128 p.
- Geyer, A. R., Smith, R. C., II, and Barnes, J. H., 1976, *Mineral collecting in Pennsylvania*: Pennsylvania Geological Survey, 4th ser., General Geology Report 33, 260 p.
- Godfrey, A. E., 1975, *Chemical and physical erosion in the South Mountain anticlinorium, Maryland*: Maryland Geological Survey, Information Circular 19, 35 p.
- Godfrey, A. E. and Cleaves, E. T., 1991, *Landscape analysis: Theoretical considerations and practical needs*: Environmental Geology and Water Sciences, v. 17, no. 2, p. 141-155.
- Goldthwait, R. P., 1976, *Frost sorted patterned ground: A review*: Quaternary Research, v. 6, p. 27-35.
- Gordon, S. E., 1922, *The mineralogy of Pennsylvania*: Philadelphia, Academy of Natural Sciences Special Publication 1, 158 p.
- Grimvall, A., Cole, C. A., Allard, B., and Sanden, P., 1986, *Quality trends of public water supplies in Sweden*: Water Quality Bulletin, v. 11, no. 4, p. 6-11.
- Hack, J. T., 1960, *Interpretation of erosional topography in humid temperate regions*: American Journal of Science Bradley Volume, v. 258-A, p. 80-97.
- Hack, J. T., 1965, *Geomorphology of the Shenandoah Valley Virginia and West Virginia and origin of the residual ore deposits*: U.S. Geological Survey Professional Paper 484, 84 p.
- Hack, J. T., 1975, *Dynamic equilibrium and landscape evolution*, in Melhorn, W. M. and Flemal, R. C., eds., *Theories of landform development*: London, George Allen and Unwin, p. 87-102.
- Haines, T. A., 1981, *Acidic precipitation and its consequences for aquatic ecosystems: a review*: Transactions of the American Fisheries Society, v. 110, p. 669-707.
- Hatcher, R. D., Jr., 1990, *Structural geology: principles, concepts, and problems*: Columbus, OH, Merrill Publishing Co., 531 p.
- Hatcher, R. D., Thomas, W. A., and Viele, G. W., 1989, *The Appalachian-Ouachita Orogen in the United States: The Geology of North America*, v. F-2: Geological Society of America, 767 p.
- Hedges, J., 1969, *Opferkessel*: Zeitschrift für Geomorphologie, v. 13, p. 22-55.
- Hedges, J., 1975, *Multiple cycles of cryoplanation on Sugarloaf Mountain, Maryland*: Biuletyn Peryglacjalny, v. 24, p. 233-243.
- Heron, S. D., Jr., Judd, J. B., and Johnson, H. S., Jr., 1971, *Clastic dikes associated with soil horizons in the North and South Carolina Coastal Plain*: Geological Society of America Bulletin, v. 82, p. 1801-1810.
- Hofmann, A. W., 1988, *Chemical differentiation of the Earth: the relationship between mantle, continental crust, and oceanic crust*: Earth and Planetary Science Letters, v. 90, p. 297-314.
- Hosterman, J. W., 1968, *White clay deposits near Mt. Holly Springs, Cumberland County, Pennsylvania*, in *The geology of mineral deposits in south-central Pennsylvania*: 33rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 38-51.

- Hosterman, J. W., 1969, White clay deposits near Mt. Holly Springs, Cumberland County, Pennsylvania: U.S. Geological Survey Professional Paper 650B, p. 66-72.
- Hupp, C. R., 1983, Geo-botanical evidence of Late Quaternary mass wasting in block field areas of Virginia: *Earth Surface Processes and Landforms*, v. 8, p. 439-450.
- Jacobson Jr., G. L., Webb III, T., and Grimm, E. C., 1987, Patterns and rates of vegetation change during the deglaciation of eastern North America, in Ruddiman W. F. and Wright, H. E., Jr., eds., *North America and adjacent oceans during the last deglaciation*: Geological Society of America, p. 277-288.
- Johnson, F., 1986, Acid precipitation: Pennsylvania Fish Commission, PFC 1003 7/86 10 M.
- Karrasch, H., 1974, Hangglättung und Kryoplanation an Beispielen aus den Alpen und kanadischen Rocky Mountains: *Akademie Der Wissenschaften, Göttingen. Mathematische-Physikale Abhandlungen*, v. 29, p. 287-300.
- Karte, J., 1982, Development and present state of German periglacial research in arctic and alpine environments: *Biuletyn Peryglacjalny*, v. 29, p. 183-201.
- Kauffman, M. E. and Frey, E. P., 1979, Antietam sandstone ridges - exhumed barrier islands or fault-bounded blocks? (abs.): *Geological Society of America Abstracts with Programs*, v. 11, no. 1, p. 18.
- Kaufmann, P. R., Herlihy, A. T., Elwood, J. W., Mitch, M. E., Overton, W. S., Sale, M. J., Messer, J. J., Cougan, K. A., Peck, D. V., Reckhow, K. H., Kinney, A. J., Christie, S. J., Brown, D. D., Hagley, C. A., and Jager, H. I., 1988, Chemical characteristics of streams in the mid-Atlantic and southeastern United States (Results of the National Stream Survey Phase I), in Vol. 1: Population descriptions and physico-chemical relationships: U.S. Environmental Protection agency, Washington, D.C., EPA/600/3-88/021a.
- Kay, M., 1951, North American Geosynclines: *Geological Society of America Memoir* 48, 143 p.
- Kemper, J., III, 1968?, American charcoal making: National Park Service, U.S. Department of the Interior, Eastern National Park and Monument Association, 25 p.
- Kimmel, W. G., Murphy, D. J., Sharpe, W. E., and DeWalle, D. R., 1985, Macroinvertebrate community structure and detritus processing rates in two southwestern Pennsylvania streams acidified by atmospheric deposition: *Hydrobiologia*, v. 124, p. 97-102.
- King, P. B., 1949, The base of the Cambrian in the southern Appalachians: *American Journal of Science*, v. 247, p. 513-530; 622-645.
- King, P. B., 1950, Geology of the Elkin area, Virginia: U.S. Geological Survey Professional Paper 230, 82 p.
- King, P. B. and Fergusen, H. W., 1960, Geology of northeasternmost Tennessee: U.S. Geological Survey Professional Paper 311, 136 p.
- Kline, S. W., Conley, J. F., and Evans, N., 1987, The Catoctin Formation in the eastern Blue Ridge of Virginia: Evidence for submarine volcanism (abs.): *Geological Society of America Abstracts with Programs*, v. 19, no. 2, p. 93.
- Kochel, R. C., 1990, Humid fans of the Appalachian Mountains, in Rachocki, A. H. and Church, M., eds., *Alluvial fans*: New York, John Wiley and Sons, p. 109-129.
- Kulander, B. R. and Dean, S. L., 1986, Structure and tectonics of the central and southern Appalachian Valley and Ridge Plateau Provinces, West Virginia, and Virginia: *American Association of Petroleum Geologists Bulletin*, v. 70, p. 1674-1684.
- Kurjack, D. C., 1958, Hopewell Village national historic site: Washington, D.C., National Park Service Historical Handbook Series No. 8, 44 p.
- Lachenbruch, A. H., 1962, Mechanics of thermal contraction cracks and ice-wedge polygons in permafrost: *Geological Society of America Special Paper* 70, 69 p.
- Laurence, R. A. and Palmer, A. R., 1963, Age of the Murray Shale and Hesse Quartzite on Chilhowee Mountain, Blount County, Tennessee: U.S. Geological Survey Professional Paper 475C, p. 53-54.
- Lauriol, B., 1990, Canadian landform examples 18: Cryoplanation terraces, northern Yukon: *Le Geographe canadien*, v. 34, no. 4, p. 347-351.
- Leffingwell, E. de K., 1915, Ground-ice wedges. The dominant form of ground-ice on the north coast of Alaska: *Journal of Geology*, v. 23, p. 635-654.
- Leffler, R. J., 1981, Estimating average temperatures on Appalachian summits: *Journal of Applied Meteorology*, v. 20, p. 637-642.
- Lesley, J. P., 1892, A summary description of the geology of Pennsylvania: Pennsylvania Geological Survey, 2nd ser., v. 1, 719 p.
- Liestøl, O., 1961, Talus taluses in Arctic regions: *Norsk Polarinstitutt Årbok*, p. 102-105.
- Lisle, R. J., 1985, Geological strain analysis, a manual for the Rf/f technique: New York, Pergamon Press, 99 p.
- Łezinski, M. W., 1909, Über die mechanische Verwitterung der Sandsteine im gemässigten Klima: *Bulletin International Akademie Sciences Crakovie, Classe Scientific Mathematique Naturelles*, v. 1, p. 1-25.
- Lynch, J. A., Corbett, E. S., and Kostelnik, K. M., 1986, Atmospheric deposition: spatial and temporal variation in Pennsylvania, 1985: Institute for Research on Land and Water Resources, The Pennsylvania State University, University Park, PA.
- Lynch, J. A., Corbett, E. S., and Kostelnik, K. M., 1987, Atmospheric deposition: spatial and temporal variation in Pennsylvania, 1986: Institute for Research on Land and Water Resources, The Pennsylvania

- State University, University Park, PA.
- McCauley, I. H., 1878, Historical sketch of Franklin County, Pennsylvania: Chambersburg, PA, D. F. Pursel, 322 p.
- Mackay, J. R., 1974, Ice-wedge cracks, Garry Island, Northwest Territories: Canadian Journal of Earth Science, v. 11, p. 1366-1383.
- Mackay, J. R., 1990, Some observations on the growth and deformation of epigenetic, syngenetic, and anti-syngenetic ice wedges: Permafrost and Periglacial Processes, v. 1, p. 15-29.
- MacLachlan, D. B. and Root, S. I., 1966, Comparative tectonics of the Cumberland and Lebanon Valleys: 31st Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, 90 p.
- McDonald, M. E., 1985, Acid deposition and drinking water: Environmental Science and Technology, v. 19, p. 772-776.
- McDonald, M. G. and Harbaugh, A. W., 1988, A modular three-dimensional finite-difference ground-water flow model: U.S. Geological Survey Techniques of Water Resources Investigations, book 6, chap. A1.
- Middlekauff, B. D., 1987, Relict periglacial morphosequences in the northern Blue Ridge: unpublished Ph.D. thesis, East Lansing, Michigan State University, 172 p.
- Mills, H. H., 1981, Boulder deposits and the retreat of mountain slopes, or, "gully gravure" revisited: Journal of Geology, v. 89, p. 649-660.
- Mills, H. H., Brakenridge, G. R., Jacobson, R. B., Newell, W. L., Pavich, M. J., and Pomeroy, J. S., 1987, Appalachian Mountains and Plateaus: Geological Society of America, Centennial Volume 2, p. 5-50.
- Mills, H. H. and Delcourt, P. A., in press, Appalachian highlands and interior low plateaus, in Morrison, R. B., ed., Quaternary non-glacial geology of the conterminous United States: Geological Society of America, Volume K-2, in press.
- Monmonier, M. S., 1967, Upland accordance in the Ridge and Valley section of Pennsylvania: unpublished M.S. thesis, University Park, The Pennsylvania State University, 58 p.
- Monmonier, M. S., 1968, Trends in upland accordance in Pennsylvania's Ridge and Valley section: Pennsylvania Academy of Science Proceedings, v. 42, p. 157-162.
- Monmonier, M. S., 1971, Upland adjustment to regional drainage in central Pennsylvania: an application of trend surface analysis: Journal of Geography, v. 70, p. 360-370.
- Mose, D. G., Diecchio, R. J., DiGiuseppi, W. H., and Nagel, M. S., 1985, Confirmation of a latest Precambrian (600 m.y.) age for the Catoctin Formation in Virginia (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 2, p. 126.
- Moss, J. H., 1976, Periglacial origin of extensive lobate colluvial deposits on the south flank of Blue Mountain near Shartlesville and Strausstown, Berks County, Pennsylvania: Pennsylvania Academy of Science Proceedings, v. 50, p. 42-44.
- Nagel, M. S. and Mose, D. G., 1984, A revised geochemical and geochronological picture of the Catoctin Formation (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 182.
- National Atmospheric Deposition Program, 1990, NADP/NTN annual data summary, precipitation chemistry in the United States, 1989.
- Neal, J. T., Langer, A. M., and Kerr, P. F., 1968, Giant desiccation polygons of Great Basin playas: Geological Society of America Bulletin, v. 79, p. 69-90.
- Nelson, A., 1968, GAF Corporation quarries--Charmian, Pennsylvania, in Cloos, E. and others, The geology of mineral deposits in south-central Pennsylvania, Annual Field Conference of Pennsylvania Geologists, 33rd, Harrisburg, PA, Guidebook, p. 6-10.
- Nickelsen, R. P., 1956, Geology of the Blue Ridge near Harpers Ferry, West Virginia: Geological Society of America Bulletin, v. 67, p. 239-270.
- Nickelsen, R. P., 1979, Sequence of structural stages of the Allegheny orogeny at the Bear Valley strip mine, Shamokin, Pennsylvania: American Journal of Science, v. 279, p. 225-271.
- Nutter, L. J., 1973, Hydrogeology of the carbonate rocks, Frederick and Hagerstown Valleys Maryland: Maryland Geological Survey, Report of Investigations No. 19, 70 p.
- Olson, C. G., 1989, Mountain soils of eastern Appalachians (abs.): 28th International Geological Congress Abstracts, v. 2, p. 2-548.
- Palmer, A. R., 1981, Subdivision of the Sauk Sequence, in Taylor, M. E., ed., Short Papers from the Second International Symposium on the Cambrian System: U.S. Geological Survey Open-File Report 81-743, p. 160-161.
- Palmer, A. R., 1983, The Decade of North American Geology, 1983 geologic time scale: Geology, v. 11, p. 503-504.
- Patterson, J. G. and Simpson, E. L., 1991, Paleoenvironmental constraints on the Chilhowee Group of the eastern Blue Ridge: Thoroughfare Gap, Virginia (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 113.
- Pavich, M. J., 1989, Regolith residence time and the concept of surface age of the Piedmont "peneplain", in Gardner, T. W. and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 181-196.
- Pearce, J. A. and Cann, J. R., 1973, Tectonic setting of basic volcanic rocks determined using trace element analysis: Earth and Planetary Science Letters, v. 19, p. 290-300.

- Pearse, J. B., 1970, A concise history of the iron manufacture of the American colonies up to the Revolution and of Pennsylvania up to the present time: New York, Burt Franklin, Research and Source Works Series 497, Selected essays in history, economics, and social sciences 142 (originally published in 1876, author "metallurgist, engineer, Commissioner of Geological Survey of Pennsylvania").
- Peltier, L. C., 1949, Pleistocene terraces of the Susquehanna River, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Bulletin G 23, 158 p.
- Peltier, L. C., 1950, The geographic cycle in periglacial regions as it is related to climatic geomorphology: Association of American Geographers Annals, v. 40, p. 214-236.
- Pettijohn, F. J., 1957, Sedimentary rocks: New York, Harper and Row, 628 p.
- Pettyjohn, W. A. and Henning, R., 1979, Preliminary estimate of ground-water recharge rates, related streamflow and water quality in Ohio: The Ohio State University Department of Geology and Mineralogy Project Completion Report No. 522, 323 p.
- Péwé, T. L., 1970, Altiplanation terraces of early Quaternary age near Fairbanks, Alaska: Acta Geographica Lodziensia, v. 24, p. 357-363.
- Péwé, T. L., 1975, Quaternary geology of Alaska: U.S. Geological Survey Professional Paper 835, 145 p.
- Péwé, T. L., 1983, The periglacial environment in North America during Wisconsin time, in Porter, S. C., ed., Late-Quaternary Environments of the United States. Volume 1. The Late Pleistocene: Minneapolis, University of Minnesota Press, p. 157-189.
- Péwé, T. L., Church, R. E., Andresen, M. J., 1969, Origin and paleoclimatic significance of large-scale patterned ground in the Donnelly Dome area, Alaska: Geological Society of America Special Paper 103, 87 p.
- Pierce, K. L., 1965, Geomorphic significance of a Cretaceous deposit in the Great Valley of southern Pennsylvania: U.S. Geological Survey Professional Paper 525-C, p. C152-C156.
- Pierce, K. L., 1966, Bedrock and surficial geology of the McConnellsburg quadrangle, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 109a, 111 p.
- Poag, C. W., and Sevon, W. D., 1989, A record of Appalachian denudation in postdrift Mesozoic and Cenozoic sedimentary deposits of the U.S. Middle Atlantic continental margin, in Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 119-157.
- Priesnitz, K., 1988, Cryoplanation, in Clark, M. J., ed., Advances in Periglacial Geomorphology: Chichester, John Wiley and Sons, p. 49-67.
- Ramsay, J. G., 1967, Folding and fracturing of rocks: New York, McGraw-Hill, 568 p.
- Ramsay, J. G. and Huber, M. I., 1983, The tectonics of modern structural geology, Vol. 1, Strain Analysis: New York, Academic Press, Inc., 307 p.
- Rankin, D. W., 1967, Guide to the geology of the Mount Rogers area, Virginia, North Carolina, and Tennessee: Carolina Geological Society Field Trip, Guidebook, 48 p.
- Rankin, D. W., 1975, The continental margin of eastern North America in the southern Appalachians: The opening and closing of the Proto-Atlantic Ocean: American Journal of Science, v. 275-A, p. 298-336.
- Rankin, D. W., Stern, T. W., Reed, J. C., Jr., and Newell, M. F., 1969, Zircon ages of felsic rocks in the Upper Precambrian of the Blue Ridge, Appalachian Mountains: Science, v. 166, p. 741-744.
- Reed, J. C., 1955, Catoctin Formation near Luray, Virginia: Geological Society of America Bulletin, v. 66, p. 871-896.
- Reger, R. D. and Péwé, T. L., 1976, Cryoplanation terraces: indicators of a permafrost environment: Quaternary Research, v. 6, p. 99-109.
- Resser, C. E. and Howell, B. J., 1938, Lower Cambrian Olenellus zone of the Appalachians: Geological Society of America Bulletin, v. 49, p. 195-248.
- Rogers, H. D., 1858, The geology of Pennsylvania: Philadelphia, v. 1, 586 p.
- Root, S. I., 1968, Geology and mineral resources of southeastern Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 119cd.
- Root, S. I., 1970, Structure of the northern terminus of the Blue Ridge in Pennsylvania: Geological Society of America Bulletin, v. 81, p. 815-830.
- Root, S. I., 1971, Geology and mineral resources of northeastern Franklin County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 119ab, 104 p.
- Root, S. I., 1977, Geology and mineral resources of the Harrisburg West area of Cumberland and York Counties, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 148ab, 106 p.
- Root, S. I., 1978, Geology and mineral resources of the Carlisle and Mechanicsburg Quadrangles, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Atlas 138ab, 1 p.
- Root, S. I., 1989, Basement control of structure in the Gettysburg rift basin, Pennsylvania and Maryland: Tectonophysics, v. 166, p. 281-292.
- Root, S. I. and Hoskins, D. M., 1977, Lat. 40°N fault-zone, Pennsylvania: a new interpretation: Geology, v. 5, p. 719-723.
- Root, S. I. and MacLachlan, D. B., 1978, Western limit of Taconic allochthons in Pennsylvania: Geological Society of America Bulletin, v. 89, p. 1515-1528.
- Rorabaugh, M. I., 1964, Estimating changes in bank storage and ground-water contribution to streamflow:

- International Association of Scientific Hydrology Publication no. 63, p. 432-441.
- Rose, A. W., 1970, Metal mines and occurrences in Pennsylvania: Pennsylvania Geological Survey, 4th ser., Mineral Resources Report 50, Part 3, 14 p.
- Russell, R. J., 1933, Alpine land forms in western United States: Geological Society of America Bulletin, v. 44, p. 927-950.
- Saad, D. A. and Hippe, D. J., 1990, Large springs in the Valley and Ridge physiographic province of Pennsylvania: U.S. Geological Survey Open-File Report 90-164, 17 p.
- Schmidlin, T. W., 1982, Leffler's method of estimating average temperatures of Appalachian summits: Evaluation in New York: Journal of Applied Meteorology, v. 21, p. 745-747.
- Schofield, C. and Trojnar, J. R., 1980, Aluminum toxicity to brook trout (*Salvelinus fontinalis*) in acidified waters, in Toribara, T. Y., Miller, M. W., and Morrow, P. E., eds., Polluted Rain: New York, Plenum Press, p. 341-365.
- Schwab, F. L., 1970, Origin of the Antietam Formation (Late Precambrian?-Lower Cambrian), central Virginia: Journal of Sedimentary Petrology, v. 40, p. 354-366.
- Schwab, F. L., 1972, The Chilhowee Group and the Late Precambrian-Early Paleozoic sedimentary framework in the central and southern Appalachians, in Lessing, P. and others, eds., Appalachian structures - origin, evolution, and possible potential for new exploration frontiers: Morgantown, West Virginia University and West Virginia Geological and Economic Survey, p. 59-101.
- Scotese, C. R. and others, 1979, Paleozoic base maps: Journal of Geology, v. 87, p. 217-277.
- Sears, C. E., 1957, Late Cretaceous erosion surface in southwest Virginia (abs.): Geological Society of America Bulletin, v. 68, p. 1883.
- Seilacher, A., 1967, Bathymetry of trace fossils: Marine Geology, v. 5, p. 413-428.
- Sepkoski, J. J., Jr. and Knoll, A. H., 1983, Precambrian-Cambrian boundary: The spike is driven and the monolith crumbles: Paleobiology, v. 9, p. 199-206.
- Sevon, W. D., 1984, A sandstone weathering rate from northeastern Pennsylvania (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 1, p. 63.
- Sevon, W. D., 1985, Pennsylvania's Polygenetic Landscape: Harrisburg Area Geological Society, 4th Annual Field Trip Guidebook, 55 p.
- Sevon, W. D., 1987, The Hickory Run boulder field, a periglacial relict, Carbon County, Pennsylvania, in Roy, D. C., ed., Northeastern Section of the Geological Society of America, Centennial Field Guide, Vol. 5: Geological Society of America DNAG Volume, p. 75-76.
- Sevon, W. D., 1989, Erosion in the Juniata drainage basin, Pennsylvania, in Gardner, T. W., and Sevon, W. D., eds., Appalachian Geomorphology (reprinted from Geomorphology, v. 2, no. 1-3): Amsterdam, Elsevier Science Publishers, p. 303-318.
- Sevon, W. D., 1990, The Hickory Run boulder field, Hickory Run State Park, Carbon County, Pennsylvania: Northeastern Geology, v. 12, p. 42-45.
- Sevon, W. D., Potter, Jr., N., and Crowl, G. H., 1983, Appalachian peneplains: an historical review: Earth Sciences History, v. 2, p. 156-164.
- Sharpe, W. E., DeWalle, D. R., Liebfried, R. T., Dinicola, R. S., Kimmel, W. G., and Sherwin, L. S., 1984, Causes of acidification of four streams on Laurel Hill in southwestern Pennsylvania: Journal of Environmental Quality, v. 13, p. 619-631.
- Shirk, W. R., 1980, Geology of southcentral Pennsylvania: Chambersburg, PA, Robson and Kaye, Inc., 136 p.
- Siddens, L. K., Seim, W. K., and Curtis, L. R., 1986, Comparison of continuous and episodic exposure to acidic, aluminum-contaminated waters of brook trout (*Salvelinus fontinalis*): Canadian Journal of Fisheries and Aquatic Sciences, v. 43, p. 2036-2040.
- Simpson, E. L., Linski, D., Mull, M. F., Keiser, J. P., Horsnall, S. L., and Hendricks, J. S., 1991, Depositional processes in outer-shelf sediments of the Lower Cambrian Harpers Formation of the Chilhowee Group, south-central Pennsylvania (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 1, p. 127.
- Simpson, E. L. and Sundberg, F. A., 1987, Early Cambrian age for synrift deposits of the Chilhowee Group of southwestern Virginia: Geology, v. 15, p. 123-126.
- Simpson, K. W., Bode, R. W., and Colquhoun, J. R., 1985, The macroinvertebrate fauna of an acid-stressed headwater stream in the Adirondack Mountains, New York: Freshwater Biology, v. 15, p. 671-681.
- Sims, S. J., 1985, Geological reconnaissance survey and reserve estimate - R. A. Bender & Son Sand Quarry, Mt. Holly Springs, Dickinson Township, Cumberland County, Pennsylvania: unpublished private report for Pennsy Supply, Inc., 18 p.
- Sloss, L. L., 1963, Sequences in the cratonic interior of North America: Geological Society of America Bulletin, v. 74, p. 93-114.
- Smith, H. T. U., and Smith, A. P., 1945, Periglacial rock streams in the Blue Ridge area (abs.): Geological Society of America Bulletin, v. 56, p. 1198.
- Smith, R. C., II, 1978, The mineralogy of Pennsylvania, 1966-1975: Friends of Mineralogy, Pennsylvania Chapter, Special Publication No. 1, 304 p.
- Smith, R. C., II and Berkheiser, S. W., Jr., 1985, Silica 99 and 44/100% pure, waiting to be mined: Pennsylvania Geology, v. 16, no. 1, p. 2-4.

- Spencer, E. W., 1988, Introduction to the structure of the Earth: New York, McGraw-Hill, 551 p.
- Spencer, E. W., Bell, J. D., and Kozak, S. J., 1989, Valley and Ridge and Blue Ridge traverse, central Virginia: 28th International Geological Congress, Washington, D.C., Field Trip Guidebook, *in* Glacial geology and geomorphology of North America, vol. 2: American Geophysical Union, 69 p.
- Stone, R. W., 1923, Roofing granules industry in southeastern Pennsylvania: Pennsylvania Geological Survey, 4th ser., Progress Report 82, 4 p.
- Stone, R. W. and Hickok, W. O., 1933, Trip 4, South Mountain, *in* Ashley, G. W., ed., 3rd Annual Field Conference of Pennsylvania Geologists, Harrisburg, PA, Guidebook, p. 11-14.
- Stose, G. W., 1906, The sedimentary rocks of South Mountain, Pennsylvania: Journal of Geology, v. 14, p. 201-220.
- Stose, G. W., 1907, White clay of South Mountain, Pennsylvania: U.S. Geological Survey Bulletin, v. 315, p. 322-334.
- Stose, G. W., 1909, Mercersburg-Chambersburg folio: U.S. Geological Survey Geologic Atlas, Folio No. 170, 19 p.
- Stose, G. W., 1932, Geology and mineral resources of Adams County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Bulletin C1, 153 p.
- Stose, G. W., 1953, Geology of the Carlisle Quadrangle, Pennsylvania: U.S. Geological Survey Geological Quadrangle Map GQ-28.
- Stose, G. W. and Bascom, F., 1929, Fairfield-Gettysburg folio: U.S. Geological Survey Geologic Atlas, Folio No. 225, 22 p.
- Stose, A. J. and Stose, G. W., 1957, Geology and mineral resources of the Gossan lead district and adjacent areas: Virginia Division of Mineral Resources Bulletin 72, 233 p.
- Swain, L. A., Hollyday, E. F., Daniel, C. C., III, Zapecza, D. S., Mesko, T. O., and Wright, W. G., 1991, Plan of study for the regional aquifer-system analysis of the Appalachian Valley and Ridge, Piedmont, and Blue Ridge physiographic provinces of the eastern and southeastern United States, with a description of study-area geology and geohydrology: U.S. Geological Survey Water-Resources Investigations Report 91-4066, 54 p.
- Swartz, F. M., 1948, Trenton and sub-Trenton of outcrop areas in New York, Pennsylvania, and Maryland: American Association of Petroleum Geologists Bulletin, v. 32, p. 1493-1595.
- Taber, S., 1943, Perennially frozen ground in Alaska: its origin and history: Geological Society of America Bulletin, v. 54, p. 1433-1548.
- Tarr, W. S., 1898, The peneplain: American Geologist, v. 21, p. 351-370.
- Taylor, F. B., Dion, J. A., and Collins, J. J., 1986, Drinking water quality and acid rain in the eastern United States: Water Quality Bulletin, v. 11, p. 50-57.
- Thomas, W. A., 1977, Evolution of the Appalachian-Ouachita salients and recesses from reentrants and promontories in the continental margin: American Journal of Science, v. 277, p. 1233-1278.
- Thornbury, W. D., 1954, Principles of Geomorphology: New York, John Wiley and Sons, 618 p.
- Thornbury, W. D., 1965, Regional geomorphology of the United States: New York, John Wiley and Sons, 609 p.
- Thornbury, W. D., 1969, Principles of Geomorphology, 2nd ed.: New York, John Wiley and Sons, 594 p.
- Tiffney, B. H., 1985, The Eocene North Atlantic land bridge: its importance in Tertiary and modern phytogeography of the northern hemisphere: Journal of the Arnold Arboretum, v. 66, p. 243-273.
- Troll, C., 1948, Der subnivale oder periglaziale Zyklus der Denudation: Erdkunde, v. 2: p. 1-21.
- Tschudy, R. H., 1965, An Upper Cretaceous deposit in the Appalachian Mountains: U.S. Geological Survey Professional Paper 525-B: p. B64-B68.
- U.S. Geological Survey, 1991, Water resources data for Pennsylvania, water year 1990--volume 2; U.S. Geological Survey Water-Data Report PA-90-2, 274 p.
- Walcott, C. D., 1891, Correlation papers, Cambrian: U.S. Geological Survey Bulletin 81, 447 p.
- Walcott, C. D., 1892a, Notes on the Cambrian rocks of Virginia and the southern Appalachians: American Journal of Science, v. 44, p. 52-57.
- Walcott, C. D., 1892b, Notes on the Cambrian rocks of Pennsylvania and Maryland from the Susquehanna to the Potomac: American Journal of Science, v. 44, p. 669-682.
- Walcott, C. D., 1893, The geologist at Blue Mountain, Maryland: National Geographic Magazine, v. 5, p. 84-88.
- Walcott, C. D., 1894a, Notes on the Cambrian rocks of Pennsylvania from the Susquehanna to the Delaware: American Journal of Science, v. 47, p. 309-311.
- Walcott, C. D., 1894b, On the Occurrence of *Olenellus* in the Green Pond Mountain series of northern New Jersey, with a note on the conglomerates: American Journal of Science, v. 47, p. 309-311.
- Walcott, C. D., 1894c, Paleozoic intraformational conglomerates: U.S. Geological Survey Bulletin 134, p. 1-43.
- Walcott, C. D., 1896, The Cambrian rocks of Pennsylvania: U.S. Geological Survey Bulletin 134, 43 p.
- Walker, J. E., 1966, Hopewell village, a social and economic history of an iron-making community: Philadelphia, University of Pennsylvania Press, 526 p.
- Wallace, P. A. W., 1961, Indians in Pennsylvania: The Pennsylvania Historical and Museum Commission, 138 p.
- Washburn, A. L., 1980, Geocryology: New York, John Wiley and Sons, 406 p.
- Washburn, A. L., 1985, Periglacial problems, *in* Church, M. and Slaymaker, O., eds., Field and Theory.

- Lectures in Geocryology: Vancouver, University of British Columbia Press, p. 166-202.
- Watts, W. A., 1979, Late Quaternary vegetation of central Appalachia and the New Jersey Coastal Plain: Ecological Monographs, v. 49, p. 427-469.
- Way, J. H., 1986, Your guide to the geology of the Kings Gap area, Cumberland County, Pennsylvania: Pennsylvania Geological Survey, 4th ser., Environmental Geology Report 8, 31 p.
- Webb, J. R., Cosby, B. J., Galloway, J. N., Hornberger, G.M., and Ryan, P. F., 1989a, The Shenandoah watershed study: An overview: Department of Environmental Sciences, University of Virginia, Charlottesville, VA.
- Webb, J. R., Cosby, B. J., Galloway, J. N., Hornberger, G.M., and Ryan, P. F., 1989b, Acidification of native brook trout streams in Virginia, a report to the Virginia Department of Game and Inland Fisheries: The Virginia Trout Stream Sensitivity Study, Project Number F-72.
- Whisonant, J. C., 1970, Paleocurrents in basal Cambrian rocks of eastern Tennessee: Geological Society of America Bulletin, v. 81, p. 2781-2786.
- Whitaker, J. C., 1955a, Direction of current flow in some Lower Cambrian clastics in Maryland: Geological Society of America Bulletin, v. 66, p. 763-766.
- Whitaker, J.C., 1955b, Geology of Catoctin Mountain, Maryland and Virginia: Geological Society of America Bulletin, v. 66, p. 435-462.
- White, S. E., 1976, Rock glaciers and block fields, review and new data: Quaternary Research, v. 6, p. 77-97.
- Wilden, R. and Mabey, D. R., 1961, Giant desiccation fissures on the Black Rock and Smoke Creek Deserts, Nevada: Science, v. 133, p. 1359-1360.
- Wilderman, C.C., 1990, Patterns of variation of pH and alkalinity in Pennsylvania streams, based on data obtained by the Alliance for Acid Rain Monitoring (ALLARM), a citizen's monitoring program, in Proceedings of the Conference on Atmospheric Deposition in Pennsylvania: A Critical Assessment, September 11-14, 1989: Environmental Resources Research Institute, The Pennsylvania State University, University Park, PA., p. 94.
- Wilderman, C. C., Cole, C. A., and Nguyen, Q. T., in press, Quality trends of some public water supplies in Pennsylvania: Northeastern Geology.
- Wilderman, C. C. and Reuss, C. L., 1991, Analysis of patterns and causes of variation in pH and alkalinity concentrations in Pennsylvania streams, using data collected by the Alliance for Acid Rain Monitoring (ALLARM): Pennsylvania Academy of Science Journal, v. 64, p. 210.
- Williams, P. J. and Smith, M. W., 1989, The Frozen Earth: Fundamentals of Geocryology: Cambridge, Cambridge University Press, Studies in Polar Research, 306 p.
- Wilshusen, J. P. and Sevon, W. D., 1981, Giant Ripples at Mount Cydonia: Pennsylvania Geology, v. 12, p. 2-8.
- Wilshusen, J. P. and Sevon, W. D., 1982, Mount Cydonia Sand Company pit, in Potter, N., Jr., ed., South Mountain: Harrisburg Area Geological Society, Guidebook, p. 28-32.
- Wolfe, J. A., 1985, Distribution of major vegetational types during the Tertiary, in Sundquist, E. T. and Broecker, W. S., eds., The carbon cycle and atmospheric CO<sub>2</sub>: natural variation Archean to present: American Geophysical Union Monograph 32, p. 357-375.
- Wood, G. D. and Clendening, J. A., 1982, Acritarchs from the Lower Cambrian Murray Shale, Chilhowee Group of Tennessee, U.S.A.: Palynology, v. 6, p. 255-265.
- Woodward, N. B., 1985, Valley and Ridge thrust belt: balanced structural sections Pennsylvania to Alabama: University of Tennessee, Department of Geological Sciences, Studies in Geology 12, 64 p.
- Worsley, P., 1984, Periglacial environment: Progress in Physical Geography, v. 8, p. 270-276.
- Zietz, I., Haworth, R. T., Williams, H., and Daniels, D. L., 1980, Magnetic anomaly map of the Appalachians: Memorial University of Newfoundland, Map No. 2, scale 1:1,000,000.