



1

NEW
INDICATION
FOR 2016

Clinical Guide to **PROBIOTIC SUPPLEMENTS**

AVAILABLE IN CANADA: 2016 Edition

Indications, Dosage Forms, and Clinical Evidence to Date

Author: Dragana Skokovic-Sunjic BScPhm RPh NCMP

Reviewers: Dr Vivien Brown MDCM CCFP FCFP NCMP, Dr Bradley C. Johnston PhD, Iris Krawchenko BScPhm RPh ACPR, Dr John Marshall MD MSc FRCPC AGAF, Dr Eamonn Quigley MD FRCP FACP MACG FRCPI, Dr Tom Smiley BScPhm PharmD

Medical Editor: Ivana Sunjic MSc

Download **PROBIOTIC GUIDE** mobile app for free



PURPOSE AND METHODS

PURPOSE:

This guide is designed to translate scientific evidence available for probiotic supplements into practical, clinically relevant information, enabling clinicians to easily select the appropriate product, dose, and format for a specific indication.

METHODS:

Published data or studies with defined clinical outcomes for probiotic strain(s) were searched, using inclusion criteria as listed in the guide. Commercially available products containing said strain(s) were identified, and the modified levels of evidence were used to rate the strength of the recommendation for each product. Once this information was compiled into a comprehensive chart form, it was assessed by independent expert reviewers.

Readers are strongly encouraged to review the evidence listed for each product for themselves.

DISCLAIMER: Information in this guide is not intended to diagnose, treat, cure, or prevent any disease. This guide is meant to be used as a clinical decision making tool to assist health care professionals in providing evidence based recommendations for their patients. In the case of probiotics, the clinical evidence supports only certain formulations/brand names of the probiotics (including the genus, species, alphanumeric designation or strain, number of live bacteria present, the blend of probiotic strains present and finally, the non-active ingredients present). Every attempt was made by the author and the reviewers to include the published clinical data for the available probiotic formulations. No bias is intended toward any specific formulation. This chart is reflective of published evidence available to date. It does not represent an endorsement of any specific product. Work on this guide is done independently and without financial conflict. In order to minimize bias, publishing and distribution efforts are coordinated via the Alliance for Education on Probiotics (AEP). We are grateful for the unrestricted publishing grant the AEP is providing for continuous efforts to translate science into clinically applicable forms.

2016 Version, valid until Dec. 31, 2016. © BHSoftInc – Please forward any questions or comments to the author Dragana Skokovic-Sunjic (dragana@bhsoftinc.com)

ACRONYMS

AAD	Antibiotic associated diarrhea - Prevention	IBD-UC	IBD - Ulcerative colitis – Adjunct to standard therapy
BG	Reduces blood glucose in Type II Diabetes (FBG and HbA1c)	IBS	Irritable bowel syndrome
C	Constipation	ID	Infectious diarrhea
CDAD	<i>Clostridium difficile</i> associated diarrhea - Prevention	LDL-C	Reduces LDL and total cholesterol
CE/AD	Childhood eczema/Atopic dermatitis	NEC*	Necrotizing Enterocolitis (newborn) *as per hospital protocol, not for self-administration
CID	Common infectious disease	NI	Nosocomial infections prevention
FAP	Functional abdominal pain	OH	Oral health 
HP	<i>Helicobacter pylori</i> - Adjunct to standard eradication therapy	Regurg/ GI Mot	Reduces regurgitation/Improves gastrointestinal motility
IBD-P	Inflammatory bowel disease - Pouchitis	TD	Traveler's diarrhea
CFU = Colony forming units (number of viable bacteria), M = million, B = billion, <i>L.</i> = <i>Lactobacillus</i> , <i>B.</i> = <i>Bifidobacterium</i>			

— INCLUSION CRITERIA AND LEVELS OF EVIDENCE

ALL 3 MUST BE SATISFIED:

1. Commercially available in Canada as a supplement or probiotic-containing food
2. GRAS - Generally Recognized as Safe status (FDA) and/or NPN - Natural Product Number (Health Canada) for probiotic strain(s) used in the products
3. Favourable published clinical evidence for the particular strain(s) present in each product/food

NOTE:

- For products containing multiple strains, evidence must be for the specified combination and not for the separate probiotic strains
- Some products are excluded from the current edition due to changes in strains used
- Please refer to individual publications for specifics of patient populations studied and details of clinical effect (age and gender)

LEVEL I: Evidence obtained from at least one properly-designed randomized trial. (HIGHEST LEVEL).

LEVEL II: Evidence obtained from well-designed controlled trials without randomization/Evidence obtained from well-designed cohort or case-control analytic studies, preferably from more than one centre or research group/Evidence obtained from multiple time series with or without the intervention. Dramatic results in uncontrolled trials might also be regarded as this type of evidence.

LEVEL III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.

INDICATIONS FOR ADULT HEALTH

Brand Name	Probiotic Strain(s)	Dosage Form	CFU/dose	No. of doses/day	ID	AAD	CDAD	TD	C	IBS	IBD-UC	IBD-P	HP	OH	LDL-C	CID
Align®	<i>B. longum</i> infantis 35624	Capsule	1B/capsule	1 capsule						¹⁻⁴						
Bio-K+® CL1285 ♦	<i>L. acidophilus</i> CL1285 <i>L. casei</i> LBC80R <i>L. rhamnosus</i> CLR2	Regular Capsule Strong Capsule "Travel Protection" Extra Strength Ferm. milk lg. Ferm. rice lg. Ferm. soy lg.	12.5B/capsule 25B/capsule 30B/capsule 50B/capsule 50B/tub 50B/tub 50B/tub	1-2 capsules 1-2 capsules 1-2 capsules 1-2 capsules 1/2 -1 tub 1/2 -1 tub 1/2 -1 tub		⁵⁻⁷	⁶⁻⁹									
BioGaia® chew tabs	<i>L. reuteri</i> DSM 17938	Chew. tab	100M/tab	1 tab	¹⁰	¹¹				¹⁶			¹²⁻¹⁵			
BioGaia® drops	<i>L. reuteri</i> DSM 17938	Drop	100M/5 drops	5 drops	¹⁰	¹¹				¹⁶			¹²⁻¹⁵			
Cardioviva™	<i>L. reuteri</i> NCIMB 30242	Capsule	2B/capsule	2 capsules												¹⁷⁻¹⁹
CulturedCare™ ProbioticGum	<i>Streptococcus salivarius</i> K12	Lozenge	0.5B/lozenge	2-10 lozenges												^{44,45}
Culturelle®	<i>L. rhamnosus</i> GG	Capsule	10B/capsule	1 capsule		²⁰							²¹			
Digestive Care™ Daily Relief	<i>L. plantarum</i> 299v	Capsule	10B/capsule	1 capsule		²²	²³			²⁴⁻²⁶						
Digestive Care™ Diarrhea Relief	<i>L. rhamnosus</i> GG	Capsule	10B/capsule	1 capsule		²⁰							²¹			
Florastor®	<i>Saccharomyces boulardii</i> lyo	Capsule Sachet	5B/capsule 5B/sachet	1-2 capsules 1-2 sachets		^{27,28}	^{8,9,28,29}	³⁰⁻³²			³³		³⁴⁻³⁶			
HMF by Genestra™	<i>L. acidophilus</i> CUL60 <i>L. acidophilus</i> CUL21 <i>B. bifidum</i> CUL20 <i>B. lactis</i> CUL34	Capsule Forte Capsule Intensive Capsule Super Powder	2.5B/capsule 10B/capsule 25B/capsule 10B/1 scoop (1 gram)	1-2 capsules 1-2 scoops			³⁷			³⁸						
IbSium®	<i>Saccharomyces cerevisiae</i> I-3856	Capsule	40B/capsule	1 capsule						³⁹						
Mutaflор® ♦	<i>Escherichia coli</i> Nissle 1917	Capsule	2.5-25B/ capsule	1-2 capsules						⁴⁰⁻⁴³						

♦ - Product requires refrigeration

INDICATIONS FOR ADULT HEALTH

Brand Name	Probiotic Strain(s)	Dosage Form	CFU/dose	No. of doses/day	ID	AAD	CDAD	TD	C	IBS	IBD-UC	IBD-P	HP	OH	LDL-C	CID
PerioBalance™	<i>L. reuteri</i> ATCC 55730 <i>L. reuteri</i> ATCC PTA 5289	100M 100M	Lozenge	200M/lozenge	2 lozenges										II ^{46,47}	
ProxiFlor® (formerly Lacidofil)	<i>L. rhamnosus</i> R0011 <i>L. helveticus</i> R0052	Capsule	4B/capsule	1-3 capsules						III ⁴⁸						
TuZen®	<i>L. plantarum</i> 299v	Capsule	10B/capsule	1-2 capsules		III ²²	III ²³			I ²⁴⁻²⁶						
Ultra Probiotic Complex by GNC	<i>L. acidophilus</i> CUL60 <i>L. acidophilus</i> CUL21 <i>B. bifidum</i> CUL20 <i>B. lactis</i> CUL34	Packet Capsule	25B/packet 25B, 50B, 75B/capsule	1-2 packets 1-2 capsules			II ³⁷			II ³⁸						
UltraFlora™ Cold Support (formerly Health Defense)	<i>L. plantarum</i> HEAL9 <i>L. paracasei</i> 8700:2	0.5B 0.5B	Capsule	1B/capsule	1 capsule											I ^{49,50}
Visbiome™ [®]	<i>L. acidophilus</i> SD5212 <i>L. casei</i> SD5218 <i>L. bulgaricus</i> SD5210 <i>L. plantarum</i> SD5209 <i>B. longum</i> SD5219 <i>B. infantis</i> SD5220 <i>B. breve</i> SD5206 <i>S. thermophilus</i> SD5207	Sachet	450B/sachet	1-2 sachets					II ⁵¹		I ⁵²⁻⁵⁴	I ^{53,55-57}				
VSL#3® [®]	<i>L. acidophilus</i> SD5212 <i>L. casei</i> SD5218 <i>L. bulgaricus</i> SD5210 <i>L. plantarum</i> SD5209 <i>B. longum</i> SD5219 <i>B. infantis</i> SD5220 <i>B. breve</i> SD5206 <i>S. thermophilus</i> SD5207	Sachet	450B/sachet	1-4 sachets					II ⁵¹		I ⁵²⁻⁵⁴	I ^{53,55-57}				

[®] - Product requires refrigeration

INDICATIONS FOR PEDIATRIC HEALTH

Brand Name	Probiotic Strain(s)	Dosage Form	CFU/dose	No. doses/day	Regurg/GI Mot	NEC* (newborn)	Colic	ID	AAD	CDAD	IBD-UC	HP	C	IBS	FAP	CID	NI	CE/AD	OH
BioGaia® chew tabs	<i>L. reuteri</i> DSM 17938	Chew. tab	100M/tab	1 tab	⁵⁸⁻⁶⁰		⁶²⁻⁶⁵	⁶⁶⁻⁷⁰	⁷¹				^{72.73}	^{74.75}	^{74.75}	^{76.77}		⁷⁸	
BioGaia® drops	<i>L. reuteri</i> DSM 17938	Drop	100M/5 drops	5 drops	⁵⁸⁻⁶⁰	⁶¹	⁶²⁻⁶⁵	⁶⁶⁻⁷⁰	⁷¹				^{72.73}	^{74.75}	^{74.75}	^{76.77}		⁷⁸	
CulturedCare™ ProbioticGum	<i>Streptococcus salivarius</i> K12	Lozenge	1B/lozenge	1-5 lozenges															^{44.45}
Culturelle®	<i>L. rhamnosus</i> GG	Powder	5B/packet	2-4 packets				⁷⁹⁻⁸⁴	^{85.86}					^{87.88}	^{87.88}		⁸⁹	⁹⁰⁻⁹⁶	
Digestive Care™ Diarrhea Relief Kids	<i>L. rhamnosus</i> GG	Powder	5B/packet	1-2 packets				⁷⁹⁻⁸⁴	^{85.86}					^{87.88}	^{87.88}		⁸⁹	⁹⁰⁻⁹⁶	
Florastor®	<i>Saccharomyces boulardii</i> lyo	Capsule Sachet	5B/capsule 5B/sachet	1-2 capsules 1-2 sachets				⁹⁸⁻¹⁰⁰	¹⁰¹	¹⁰²		¹⁰⁵							
HMF Fit for School™ [50 mg vitamin C]	<i>L. acidophilus</i> CUL21 <i>L. acidophilus</i> CUL60 <i>B. lactis</i> CUL34 <i>B. bifidum</i> CUL20	Tablet	12.5B/tablet	1 tablet													¹⁰³		
Junior Daily Probiotics (formerly DDS® Junior)	<i>B. lactis</i> UABLA-12 4.2B <i>L. acidophilus</i> DDS®-1 0.8B	Powder	5B/gram	2 grams												¹⁰⁴		¹⁰⁵	
Probiotic Baby	<i>B. lactis</i> BB-12	Drop	1B/6 drops	6 drops					¹⁰⁷							¹⁰⁶			
Proxiflor® (formerly Lacidofil)	<i>L. rhamnosus</i> R0011 <i>L. helveticus</i> R0052	Capsule	4B/capsule	1 capsule														¹⁰⁸	

* as per hospital protocol, not for self-administration

INDICATIONS FOR PEDIATRIC HEALTH

Brand Name	Probiotic Strain(s)	Dosage Form	CFU/dose	No. doses/day	Regurg/GI Mot	NEC* (newborn)	Colic	ID	AAD	CDAD	IBD-UC	HP	C	IBS	FAP	CID	NI	CE/AD	OH
Ultimate FLORA™ [®] (formerly FloraBABY)	<i>B. breve</i> HA-129 <i>L. rhamnosus</i> HA-111 <i>B. bifidum</i> HA-132 <i>B. infantis</i> HA-116 <i>B. longum</i> HA-135	1.2B 1B 0.8B 0.6B 0.4B	Powder	4B/scoop	1 scoop	II ⁹⁷													
UltraFlora™ Baby	<i>L. rhamnosus</i> GG <i>B. lactis</i> BB-12	Drop	1B/6 drops	6 drops												II ¹⁰⁹			
UltraFlora™ Children's	<i>L.acidophilus</i> NCFM® <i>B.animalis</i> subsp. <i>lactis</i> Bi-07	2.5B 2.5B	Chewable tablet	5B/chew. tablet	1-2 chew. tablets											I ¹¹⁰			
Visbiome™ [®]	<i>L.acidophilus</i> SD5212 <i>L.casei</i> SD5218 <i>L.bulgaricus</i> SD5210 <i>L.plantarum</i> SD5209 <i>B.longum</i> SD5219 <i>B.infantis</i> SD5220 <i>B.breve</i> SD5206 <i>S.thermophilus</i> SD5207		Sachet	450B/sachet	1-2 sachets			I ¹¹¹			I ^{112,113}		I ¹¹⁴						
VSL#3® [®]	<i>L.acidophilus</i> SD5212 <i>L.casei</i> SD5218 <i>L.bulgaricus</i> SD5210 <i>L.plantarum</i> SD5209 <i>B.longum</i> SD5219 <i>B.infantis</i> SD5220 <i>B.breve</i> SD5206 <i>S.thermophilus</i> SD5207		Sachet	450B/sachet	1-2 sachets			I ¹¹¹			I ^{112,113}		I ¹¹⁴						

[®] - Product requires refrigeration

* as per hospital protocol, not for self-administration

INDICATIONS FOR VAGINAL HEALTH

Brand Name	Probiotic Strain(s)	Dosage Form	CFU per dose	No. of doses/day	Vulvovaginal candidiasis	Bacterial vaginosis	
Fem-Dophilus®	<i>L. rhamnosus</i> GR-1 <i>L. reuteri</i> RC-14	2.5B 2.5B	Oral capsule	5B/capsule	1 capsule	[115,116]	[117-119]
ProB™ (RePhresh ProB)	<i>L. rhamnosus</i> GR-1 <i>L. reuteri</i> RC-14	2.5B 2.5B	Oral capsule	5B/capsule	1 capsule	[115,116]	[117-119]
Probaclic BV®	<i>L. acidophilus</i> A-212 <i>L. rhamnosus</i> A-119 <i>S. thermophilus</i> A-336	0.4B 6.8B 0.8B	Vaginal capsule	8B/capsule	1-2 capsules		[120]
Provacare™	<i>L. rhamnosus</i> Lcr35		Vaginal capsule	3.41B/capsule	2 capsules	[121]	[122-124]
Purfem™	<i>L. rhamnosus</i> PB01 <i>L. gasseri</i> EN-153471 (EB01)	1B 1B	Vaginal ovule	2B/ovule	1 ovule		[125,126]
UltraFlora™ Women's	<i>L. reuteri</i> RC-14 <i>L. rhamnosus</i> GR-1	1B 1B	Oral capsule	2B/capsule	2 capsules	[127]	[128,129]

FUNCTIONAL FOODS WITH ADDED PROBIOTICS

Brand Name	Probiotic Strain(s)	CFU/dose	No. of doses / day	Children				Adults				
				ID	HP	AAD	CID	AAD	IBS	HP	C	BG
Activia® *	<i>B. (animalis) lactis</i> CNCM I-2494	1B/serving	1-3 servings						130,131		132	
DanActive® *	<i>L. casei</i> sp. <i>Paracasei</i> CNCM I-1518	10B/serving	1-2 servings	133-135	136		137,139,140	138				141-148
iOGO ProBio™ *	<i>B. lactis</i> BB-12 <i>L. acidophilus</i> LA-5	1B/100g	1-3 servings							150-153	154,155	156
Nestlé® Gerber® Graduates Infant Cereals	<i>B. lactis</i> BB-12 DSM 10140	1B/28g serving	1 serving			106,149	106,149					
Nestlé® Gerber® Graduates Toddler Cereals	<i>B. lactis</i> BB-12 DSM 10140	1B/35g serving	1 serving			106,149	106,149					
Nestlé® Gerber® Graduates Toddler Drink	<i>B. lactis</i> BB-12 DSM 10140	1B/200mL serving	1 serving			106,149	106,149					
Nestlé® Good Start® Probiotic Infant Formula	<i>B. lactis</i> BB-12 DSM 10140	130M/100mL serving	Routine feeding if an alternative to breast milk is required			107						
Yoptimal® *	<i>B. lactis</i> BB-12 <i>L. acidophilus</i> LA-5	1B/100g	1-3 servings							150-153	154,155	156

* - Product requires refrigeration

REFERENCES

1. Brenner DM, Chey WD. *Bifidobacterium infantis* 35624: A novel probiotic for the treatment of IBS. *Rev Gastroenterol Disord.* 2009;9(1):7-15.
2. Quigley EM, Whorwell PJ, et al. Probiotic use results in normalization of bowel movement frequency in IBS, results from a clinical trial with the novel probiotic *Bifidobacterium infantis* 35624. *Am J Gastroenterol.* 2005;100(9 Suppl A):S26.
3. Whorwell PJ, Altrichter L, et al. Efficacy of an encapsulated probiotic *Bifidobacterium infantis* 35624 in women with irritable bowel syndrome. *Am J Gastroenterol.* 2006;101(7):1581-90.
4. O'Mahony L, McCarthy J, Kelly P, Hurley G, Luo F, Chen K, O'Sullivan GC, Kiely B, Collins JK, Shanahan F, Quigley EM. Gastroenterology. 2005;128:541-51. *Lactobacillus* and *bifidobacterium* in irritable bowel syndrome: symptom responses and relationship to cytokine profiles.
5. Beausoleil M, Fortier N, et al. Effect of a fermented milk combining *Lactobacillus acidophilus* CL1285 and *Lactobacillus casei* in the prevention of antibiotic-associated diarrhea: A randomized, double-blind, placebo-controlled trial. *Can J G*
6. Gao XW, et al. Dose-response efficacy of a proprietary probiotic formula of *Lactobacillus acidophilus* CL1285 and *Lactobacillus casei* LBC80R for antibiotic-associated diarrhea and clostridium difficile-associated diarrhea prophylaxis in adult
7. Sampalis J, Psaradellis E, Rampakakis E. Efficacy of BIO K+ CL1285 in the reduction of antibiotic-associated diarrhea - a placebo controlled double-blind randomized, multi-center study. *Arch Med Sci* 2010; 6: 56-64.
8. Johnston BC, Ma SS, et al. Probiotics for the prevention of Clostridium difficile- associated diarrhea: A systematic review and meta-analysis. *Ann Intern Med.* 2012.
9. Johnson S, Maziade P, et al. Is primary prevention of Clostridium difficile infection possible with specific probiotics? *Int J Infect Dis.* 2012;16(11):e786-92.
10. Aiello R, Ali Mea. Efficacy and safety of *Lactobacillus reuteri* ATCC 55730 in preventing acute chemotherapy induced diarrhoea in colon cancer patients treated with DeGramont or folfox schedule. *Ann Oncol.* 2008;19(suppl 9):abstract B7.
11. Cimperman L, Best K, et al. A randomized, double-blind, placebo-controlled pilot study of *Lactobacillus reuteri* for the prevention of antibiotic-associated diarrhea in hospitalized adults. *J Parenter Enteral Nutr.* 2009;33(229):abstract SP-31.
12. Ainora M, Nista EC, et al. Impact of *L. reuteri* supplementation on anti-*H. pylori* second line therapy-related. *Gut.* 2008;57 Suppl II(A327): abstract P1111.
13. Francavilla R, Lionetti E, et al. Inhibition of *Helicobacter pylori* infection in humans by *Lactobacillus reuteri* ATCC55730 and effect on eradication therapy: A pilot study. *Helicobacter.* 2008;13:127-34.
14. Imae K, Tanaka A, et al. *Lactobacillus reuteri* tablets suppress *Helicobacter pylori* infection -a double-blind, randomized, placebo-controlled cross-over clinical study. *J Jap Assoc Inf Dis.* 2007;81:387-93.
15. Saggiorno A, Caroli M, et al. *Helicobacter pylori* eradication with *Lactobacillus reuteri*. A double blind placebo-controlled study. *Dig Liver Dis.* 2005;37(Suppl 1):S88.
16. Ojetti, V., Ianiro, G., Tortora, A., D'Angelo, G., Di Renzo, T. A., Bibbo, S., Migneco, A., Gasbarrini, A. The effect of *Lactobacillus reuteri* supplementation in adults with chronic functional constipation: a randomized, double-blind, placebo- controlled trial. 2014
17. Jones ML, Martoni C, Prakash S. Cholesterol lowering and inhibition of sterol absorption by *Lactobacillus reuteri* NCIMB 30242: a randomized controlled trial. *European Journal of Clinical Nutrition.* 2012 Nov;66(11):1234-41.
18. Jones ML, Martoni C, Parent M, Prakash S. Cholesterol lowering efficacy of a microencapsulated BSH-active *Lactobacillus reuteri* NCIMB 30242 yoghurt formulation in hypercholesterolemic adults. *British Journal of Nutrition.* 2012 May;107(10):15
19. Jones ML, Tomaro-Duchesneau C, Martoni C, Prakash S. Cholesterol-lowering with Bile Salt Hydrolase (BSH) active probiotic bacteria, mechanism of action, clinical evidence and future direction for heart health applications. *Expert Opinion on*
20. Kale-Pradhan PB, Jassal HK, Wilhelm SM. The use of *Lactobacillus* in the prevention of antibiotic associated diarrhea: A meta-analysis. *Pharmacotherapy.* 2010;30(2):119-26.
21. Armuzzi A, Cremonini F, et al. The effect of oral administration of *Lactobacillus GG* on antibiotic-associated gastrointestinal side effects during *Helicobacter pylori* eradication therapy. *Alimentary Pharmacology & Therapeutics.* 2001;15(2):163
22. Lonnermark E, Friman V, et al. Intake of *Lactobacillus plantarum* reduces certain gastrointestinal symptoms during treatment with antibiotics. *J Clin Gastroenterol.* 2010;44(2):106-12.
23. Klarin B, Wullt M, et al. *Lactobacillus plantarum* 299v reduces colonisation of *Clostridium difficile* in critically ill patients treated with antibiotics. *Acta Anaesthesiologica Scandinavica.* 2008;59:1096-102.
24. Niedzielin K, Korodecki H, Birkenfeld B. A controlled, double-blind, randomized study on the efficacy of *Lactobacillus plantarum* 299V in patients with irritable bowel syndrome. *Eur J Gastroenterol Hepatol.* 2001;13(10):1143-7.
25. Nobaek S, Johansson ML, et al. Alteration of intestinal microflora is associated with reduction in abdominal bloating and pain in patients with irritable bowel syndrome. *Am J Gastroenterol.* 2000;95:1231.
26. Ducrotte P, Sawant PD, Jayanthi V. Clinical trial: *Lactobacillus plantarum* 299v (DSM 9843) improves symptoms of irritable bowel syndrome. *World J Gastroenterol.* 2002;18(30):4012-8.
27. Can M, Besirbileoglu BA, et al. *Prophylactic Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea: A prospective study. *Med Sci Monit.* 2006;12(4):19-22.
28. Surawicz CM, McFarlane LV, et al. The search for a better treatment for recurrent Clostridium difficile disease: Use of high-dose vancomycin combined with *Saccharomyces boulardii*. *Clin Infect Dis.* 2000;31:1012-7.
29. McFarlane LV, Surawicz CM, et al. A randomized placebo-controlled trial of *Saccharomyces boulardii* in combination with standard antibiotics for Clostridium difficile disease. *JAMA.* 1994;271:1913-8.
30. Kollaritsch HH, Kremsner P, et al. Prevention of traveler's diarrhea: Comparison of different non-antibiotic preparations. *Travel Medicine International.* 1989;1:9.
31. Kollaritsch HH, et al. Prophylaxis of traveller's diarrhea with *Saccharomyces boulardii*, results of a placebo-controlled double blind study. *Fortschr Med.* 1993;111(9):152-6.
32. McFarlane LV. Meta-analysis of probiotics for the prevention of traveler's diarrhea. *Travel Med Infect Dis.* 2007;5:97-105.
33. Guslardi M, Mezzi G, et al. *Saccharomyces boulardii* in maintenance treatment of Crohn's disease. *Dig Dis Sci.* 2000;45(7):1462-4.
34. Song MJ, Park DI, et al. The effect of probiotics and mucoprotective agents on PPI-based triple therapy for eradication of *Helicobacter pylori*. *Helicobacter.* 2010;15(3):206-13.
35. Szajewska H, Horvath A, Piwowarczyk A. Meta-analysis: The effects of *Saccharomyces pylori* eradication rates and side effects during treatment. *Alimentary Pharmacology & Therapeutics.* 2010;32(9):1069-7.
36. Cindoruk M, Erkan G, et al. Efficacy and safety of *Saccharomyces boulardii* in the 14-day triple anti-*Helicobacter pylori* therapy: a prospective randomized placebo-controlled double-blind study. *Helicobacter.* 2007;12: 309-16.
37. Plummer S, et al. 2004. Clostridium difficile pilot study: effects of probiotic supplementation on the incidence of C. difficile diarrhea. *International Microbiol.* 7, 59-62
38. Williams E. A., et al. 2008. Clinical trial: a multistrain probiotic preparation significantly reduces symptoms of irritable bowel syndrome in a double-blind placebo-controlled study. *Aliment Pharmacol Ther.* 29, 97-103

39. Pineton de Chambrun, G., Neut, C., Chau, A., Cazaubiel, M., Pelerin, F., Justen, P., and Desreumaux, P. (2015). A randomized clinical trial of *Saccharomyces cerevisiae* versus placebo in the irritable bowel syndrome. *Digestive And Liver Disease* 47, 119-124.
40. Floch MH. 1. editorials.doi:10.1097/MCG.0b013e3181cf83f7. *Journal of Clinical Gastroenterology*. 2010;44(4):237-8.
41. Henker J, Muller S, et al. Probiotic Escherichia coli Nissle 1917 (EcN) for successful remission maintenance of ulcerative colitis in children and adolescents: An open-label pilot study. *Zeitschrift fuer Gastroenterologie*. 2008;46(9):874-5.
42. Kruis W, et al. Double-blind comparison of an oral Escherichia coli preparation and mesalazine in maintaining remission of ulcerative colitis. *Alimentary Pharmacology & Therapeutics*. 1997;11:853-8.
43. Kruis W, et al. Maintaining remission of ulcerative colitis with the probiotic Escherichia coli Nissle 1917 is as effective as with standard mesalazine. *Gut*. 2004;53:167-23.
44. Burton, J.P., Chilcott, C.N., Moore, C.J., Speiser, G., Tagg, J.R. A preliminary study of the effect of probiotic *Streptococcus salivarius* K12 on oral malodour parameters." *J App Microb*.
45. Di Pierro, F., Donato, G., Fomia, F., Adamo, T., Careddu, D., Cassandro, C., Albera, R. "Preliminary pediatric clinical evaluation of the oral probiotic Streptococcus salivarius K12 in preventing recurrent pharyngitis and/or tonsillitis caused by *Streptococcus pyogenes* and recurrent acute otitis media." *2012 Int J Gen Med*
46. Vivekananda MR, Vandana KL, Bhar KG. Effect of the probiotic Lactobacillus reuteri (Prodentis) in the management of periodontal disease: a preliminary randomized clinical trial. *Journal of Oral Microbiology* 2010, 2: 5344 - DOI: 10.3402/jom.v
47. Twetman S et al. Short-term effect of chewing gums containing probiotic *Lactobacillus reuteri* on the levels of inflammatory mediators in gingival crevicular fluid. *Acta Odontol Scand*. 67(1):19-24.
48. Benes, Z, et al. A probiotic combination for IBS: a pilot clinical study. *NUTRAFoods* : 2006, 5: 20-27.
49. Berggren A et al. Randomised, double-blind and placebo-controlled study using new probiotic *lactobacilli* for strengthening the body immune defence against viral infections; *Eur J Nutr*. 2011;50(3):203-210
50. Busch R et al. Randomized, Double Blind and Placebo Controlled Study Using a Combination of Two Probiotic *Lactobacilli* to Alleviate Symptoms and Frequency of Common Cold. *Food and Nutrition Sciences*, 2013, 4(1A), 13-20
51. Parkes GC, Chattoor D, Emmanuel A. The probiotic VSL#3 increases scbm and reduces symptom severity scores in patients with functional constipation. *Gut*. 2011;60:163.
52. Sood A, Midha V, et al. The probiotic preparation VSL#3 induces remission in patients with mild-to-moderately active ulcerative colitis. *Clin Gastroenterol Hepatol*. 2009;11:1202-9.
53. Tursi A, Brandimarte G, et al. Low-dose balsalazide plus a high-potency probiotic preparation is more effective than balsalazide alone or mesalazine in the treatment of acute mild-to-moderate ulcerative colitis. *Med Sci Monit*. 2004;10(11):PI
54. Tursi A, Brandimarte G, et al. Treatment of relapsing mild-to-moderate ulcerative colitis with the probiotic VSL#3 as adjunctive to a standard pharmaceutical treatment: A double-blind, randomized, placebo-controlled study. *Am J Gastroenterol*
55. Gionchetti P, et al. Oral bacteriotherapy as maintenance treatment in patients with chronic pouchitis: A double-blind, placebo-controlled trial. *JGastro*. 2000;119(2):305-9. *Gastro*. 2000;119(2):305-9.
56. Gionchetti P, et al. Prophylaxis of pouchitis onset with probiotic therapy: A double-blind, placebo-controlled trial. *Gastroenterology*. 2003;124(5):1202-9.
57. Mimura T, Rizzello F, et al. Once daily high dose probiotic therapy (VSL#3) for maintaining remission in recurrent or refractory pouchitis. *Gut*. 2004;53:108-14.
58. Indrio F, Riezzo G, et al. The effect of probiotics on feeding tolerance, bowel habits, and gastrointestinal motility in preterm newborns, a double-blind, placebo-controlled pilot study. *J Pediatr*. 2008;152:801-6.
59. Indrio F, Riezzo G, et al. *Lactobacillus reuteri* accelerates gastric emptying and improves regurgitation in infants. *Eur J Clin Invest*. 2012;41(4):417-22.
60. Garofoli F, Civardi E, et al. The early administration of *Lactobacillus reuteri* DSM 17938 controls regurgitation episodes in full-term breastfed infants. *Int J Food Sci Nutr*. 2014
61. Hunter et al. Effect of routine probiotic, *Lactobacillus reuteri* DSM 17938, use on rates of necrotizing enterocolitis in neonates with birthweight < 1000 grams: a sequential analysis , *BMC Pediatrics* 2012, 12:142
62. Savino F, Pelle E, Palumeri E, et al. *Lactobacillus reuteri* (ATCC 55730) versus simethicone in the treatment of infantile colic: A prospective randomized study. *Pediatrics*. 2007;119:124-30.
63. Savino F, Cordisco L, et al. *Lactobacillus reuteri* DSM 17 938 in infantile colic: A randomized, double-blind, placebo-controlled study. *Pediatrics*. 2010;126(3):e526-533.
64. Sung V, Hiscock H, et al. Treating infant colic with the probiotic *Lactobacillus reuteri*: double blind, placebo controlled randomised trial. *BMJ*. 2014;1(348):g2107.
65. Chau K, et al. Probiotics for Infantile Colic: A Randomized, Double-Blind, Placebo-Controlled Trial Investigating *Lactobacillus reuteri* DSM 17938; *J Pediatrics* January 2015 Volume 166, Issue 1, Pages 74-78.e1 Published Online: October 22, 2014
66. Eom TH, Oh EY, et al. The therapeutic effect of *Lactobacillus reuteri* in acute diarrhea in infants and toddlers : A prospective, randomized, placebo-controlled study. *Korean J Ped*. 2005;48:986-9.
67. Shornikova AV, Casa IA, et al. *Lactobacillus reuteri* as a therapeutic agent in acute diarrhea in young children: A prospective, randomized, placebo-controlled study. *J Pediatr Gastroenterol Nutr*. 1997;24:399-404.
68. Agustina R, Kok FJ, et al. Randomized trial of probiotics and calcium on diarrhea and respiratory tract infections in Indonesian children. *Pediatrics*. 2012;95(5):1-10.
69. Francavilla R, Lionetti E, et al. Randomised clinical trial: *Lactobacillus reuteri* DSM 17938 vs. placebo in children with acute diarrhoea - a double-blind study. *Alimentary Pharmacology & Therapeutics*. 2012;36(4):363-9.
70. Dinleyici EC, PROBAGE Study Group. Vandenplas Y. *Lactobacillus reuteri* DSM 17938 effectively reduces the duration of acute diarrhoea in hospitalised children. *Acta Paediatr*. 2014
71. Lionetti E, Minnelli VL, et al. *Lactobacillus reuteri* therapy to reduce side effects during anti-*Helicobacter pylori* treatment in children: A randomized placebo controlled trial. *Alimentary Pharmacology & Therapeutics*. 2006;24:1461-8.
72. Coccorullo P, Strisciuglio P, et al. *Lactobacillus reuteri* (DSM 17938) in infants with functional chronic constipation: A double-blind, randomized, placebo-controlled study. *J Pediatr*. 2010;157(4):598-602.
73. Indrio F, Di Mauro A, et al. Prophylactic use of a probiotic in the prevention of colic, regurgitation, and functional constipation: a randomized clinical trial. *JAMA Pediatr*. 2013;168(3):228-233.
74. Romano C, Ferrau V, et al. *Lactobacillus reuteri* in children with functional abdominal pain (FAP). *J Paediatr Child Health*. 2010 2010 Jul 8.
75. Weizman Z, et al. A Randomized Controlled Trial of *Lactobacillus reuteri* DSM 17938 in Functional abdominal pain of childhood, SP-N-0102, Nutrition Trial symposia within the ESPGHAN program (9-12th of June, 2014 presented at 47th Annual Meet
76. Weizman Z, Asli G, Alsheikh A. Effect of a probiotic infant formula on infections in child care centers: Comparison of two probiotic agents. *Pediatrics*. 2005;115(1):5-9.
77. Gutierrez- Castrellon P, Lopez- Velazquez Get al. Diarrhea in preschool children and *Lactobacillus reuteri*: a randomized controlled trial. *Pediatrics*. 2014;133:e904.
78. Abrahamsson TR, Jakobsson T, et al. Probiotics in prevention of IgE associated eczema: a double-blind, randomized, placebo-controlled trial. *J Allergy Clin Immunol* 2007; 119: 1174-1180.
79. Szajewska H, Mrukowicz JZ. Probiotics in the treatment and prevention of acute infectious diarrhea in infants and children: A systematic review of published randomized, double-blind, placebo-controlled trials. *J Pediatr Gastroenterol Nutr*. 2

80. Basu S, Chatterjee M, Ganguly S, Chandra P.K. Efficacy of *Lactobacillus rhamnosus* GG in acute watery diarrhea of Indian children: a randomised controlled trial. *Journal of Paediatrics and Child Health*. 2007;43(12):837-42.
81. Basu S, Paul DK, et al. Efficacy of high-dose *Lactobacillus rhamnosus* GG in controlling acute watery diarrhea in Indian children: a randomized controlled trial. *Journal of Clinical Gastroenterology* 2009;43(3):208-13.
82. Guandalini S, Pensabene L, et al. *Lactobacillus* GG administered in oral rehydration solution to children with acute diarrhoea: a multicenter European trial. *Journal of Pediatric Gastroenterology and Nutrition* 2000;30(1):54-60.
83. Ritchie BK, Brewster DR, et al. Efficacy of *Lactobacillus* GG in Aboriginal children with acute diarrhoeal disease: a randomised clinical trial. *Journal of Pediatric Gastroenterology and Nutrition* 2010;50(6):619-24.
84. Misra S, Sabui TK, Pal NK. A randomized controlled trial to evaluate the efficacy of *Lactobacillus* GG in infantile diarrhea. *The Journal of Pediatrics* 2009;155(1):129-32.
85. Vanderhoof JA, Whitney DB, et al. *Lactobacillus* GG in the prevention of antibiotic-associated diarrhea in children. *J Pediatr*. 1999;135:564-8.
86. Johnston BC, Goldenberg JZ, et al. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database Syst Rev*. 2011;11:CD004827.
87. Gawronski A, Dziechciarz P, et al. A randomized, double-blind placebo-controlled trial of *Lactobacillus* GG for abdominal pain disorders in children. *Alimentary Pharmacology & Therapeutics*. 2007;25(2):177-84.
88. Francavilla R, et al. A Randomized Controlled Trial of *Lactobacillus* GG in Children With Functional Abdominal Pain. *Pediatrics* 2010; 126:e1445-e1452
89. Hojsak I, et al. *Lactobacillus* GG in the Prevention of Nosocomial Gastrointestinal and Respiratory Tract Infections. *Pediatrics* 2010; 125:e1171-e1177;
90. Doege K, Grajecki D, et al. Impact of maternal supplementation with probiotics during pregnancy on atopic eczema in childhood—a meta-analysis. *Br J Nutr*. 2012;107:1-6.
91. Kalliomaki M, Salminen S, et al. Probiotics in primary prevention of atopic disease: a randomised placebo controlled trial. *Lancet*. 2001;357:1076-1079.
92. Rautava S, Kalliomaki M, Isolauri E. Probiotics during pregnancy and breast-feeding might confer immunomodulatory protection against atopic disease in the infant. *J Allergy Clin Immunol*. 2002;109:119-121.
93. Kalliomaki M, Salminen S, et al. Probiotics and prevention of atopic disease: 4-year follow-up of a randomised placebo-controlled trial. *Lancet*. 2003;361:1869-1871.
94. Kalliomaki M, Salminen S, et al. Probiotics during the first 7 years of life: a cumulative risk reduction of eczema in a randomized, placebo controlled trial. *J Allergy Clin Immunol*. 2007;119:1019-1021.
95. Dotterud C, Større O, et al. Probiotics in pregnant women to prevent allergic disease: a randomized, double-blind trial. *Br J Dermatol* 2010; 163: 616-623
96. Boyle RJ, Ismail IH, et al. *Lactobacillus* GG treatment during pregnancy for the prevention of eczema: a randomized controlled trial. *Allergy* 2011; 66: 509-516.
97. Janvier A, Malo J, Barrington K. Cohort study of probiotics in a North American neonatal intensive care unit. *J Pediatr* 2014;164:980-5
98. Htwe Y, Kee KS, et al. Effect of *Saccharomyces boulardii* in the treatment of acute watery diarrhea in Myanmar children: A randomized controlled study. *Am J Trop Med Hyg*. 2008;78:214-6.
99. Kurugol Z, Koturoglu G. Effects of *Saccharomyces boulardii* in children with acute diarrhea. *Acta Paediatrica*. 2005;94:44-7.
100. Feizizadeh S, Salehi-Abargouei A, Akbar V Efficacy and Safety of *Saccharomyces boulardii* for Acute Diarrhea . *Pediatrics* 2014; 134:e176-e191
101. Kotowska M, Albrecht P, Szajewska H. *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea in children: A randomized double-blind placebo-controlled trial. *Alimentary Pharmacology & Therapeutics*. 2005;21(5):583-90.
102. Buts JP, Corthier G, Delmee M. *Saccharomyces boulardii* for Clostridium difficile-associated enteropathies in infants. *J Pediatr Gastroenterol Nutr*. 1993;16:419-25.
103. Garaiova I, Muchová J, Nagyová Z, Wang D, Li JV, Országová Z, Michael DR, Plummer SF, Duracková Z. (2015) Probiotics and vitamin C for the prevention of respiratory tract infections in children attending preschool: a randomised controlled pilot study. *Eur J Clin Nutr*. 69(3):373-9
104. Gerasimov S, et al. Role of probiotics in attenuation of acute respiratory tract infections in preschool and primary school children, Abstract Group: 74. Paediatric Respiratory Infection and Immunology. Abstract Number : 1641 at European Re
105. Gerasimov S. Probiotic supplement reduces atopic dermatitis in preschool children: a randomized, double-blind, placebo-controlled, clinical trial. *Am J Clin Dermatol*. 2010; 11(5):351-61.
106. Taipale T, Pienihäkinen K, et al. *Bifidobacterium animalis* subsp. *lactis* BB-12 in reducing the risk of infections in infancy. *Br J Nutr*. 2011;105:409-416.
107. Corréa NBO, Filho LAP, et al. A Randomized Formula Controlled Trial of *Bifidobacterium lactis* and *Streptococcus thermophilus* for Prevention of Antibiotic-Associated Diarrhea in Infants. *J Clin Gastroenterol*. 2005;39(5):385-389.
108. Chernyshov, P. Integrated treatment of infants, patients with atopic dermatitis. [Dermatology] 2007, 3: 23-26.
109. Rautava S, Salminen S, Isolauri E. Specific probiotics in reducing the risk of acute infections in infancy—a randomised, double-blind, placebo-controlled study. *Br J Nutr*. 2009;101:1722-1726
110. Layer G, et al. Probiotics effect on cold and influenza-like symptoms incidence and duration in children; *J Pediatrics* 2009; 124:e172-e179
111. Dubey AP, et al. Use of VSL#3® (a new high concentration probiotic mixture) in the treatment of childhood diarrhea with specific reference to rotavirus diarrhea. *J Clin Gastroenterol*. 2008;42(Suppl 3 Pt 1):S126-9.
112. Huynh HQ, deBruyn J, et al. Probiotic preparation VSL#3 induces remission in children with mild to moderate acute ulcerative colitis: A pilot study. *Inflamm Bowel Dis*. 2009;15:760-8.
113. Miele E, Pascarella F, et al. Effect of a probiotic preparation (VSL#3) on induction and maintenance of remission in children with ulcerative colitis. *Am J Gastroenterol*. 2009;104:437-43.
114. Guandalini S, Magazzù G, et al. VSL#3 improves symptoms in children with irritable bowel syndrome: A multicenter, randomized, placebo-controlled, double-blind, crossover study. *J Pediatr Gastroenterol Nutr*. 2012;51(1):24-30
115. Anukam K, Osazuwa E, et al. Augmentation of antimicrobial metronidazole therapy of bacterial vaginosis with oral probiotic *Lactobacillus rhamnosus* GR-1 and *Lactobacillus reuteri* RC-14: Randomized, double-blind, placebo controlled trial. *Micr*.
116. Reid G, Charbonneau D, et al. Oral use of *Lactobacillus rhamnosus* GR-1 and *L. fermentum* RC-14 significantly alters vaginal flora: Randomized, placebo-controlled trial in 64 healthy women. *FEMS Immunol Med Microbiol*. 2003;35:131-4.
117. Anukam K, Osazuwa E, et al. Clinical study comparing probiotic *Lactobacillus* GR-1 and RC-14 with metronidazole vaginal gel to treat symptomatic bacterial vaginosis. *Microbes and Infection*. 2006;8(12-13):2772-6.
118. Hummelen R, Changalucha J, et al. *Lactobacillus rhamnosus* GR-1 and *L. reuteri* RC-14 to prevent or cure bacterial vaginosis among women with HIV. *Int J Gynaecol Obstet*. 2010;111(3):254-8.
119. Reid G, Bruce AW, et al. Oral probiotics can resolve urogenital infections. *FEMS Immunol Med Microbiol*. 2001;30:49-52.
120. Ya W, Reifer C, Miller LE. Efficacy of vaginal probiotic capsules for recurrent bacterial vaginosis: a double-blind, randomized, placebo-controlled study. *American journal of obstetrics and gynecology*, 2010. 203 (2), pp. 120-1.
121. Kern AM, Bohbot JM, Cardot JM. Preventive treatment of vulvovaginal candidosis with vaginal probiotic (gynophilus®-lcr regenerans®): Results of the observational study candiflore. *La Lettre du Gynécologue*. 2012;370:33-7.

122. Marcone V, Rocca G, et al. Long-term vaginal administration of *Lactobacillus rhamnosus* as a complementary approach to management of bacterial vaginosis. *Int J Gynaecol Obstet.* 2010;110(3):223-6.
123. Petricovic L, Witt A. The role of *Lactobacillus casei/rhamnosus* Lcr35 in restoring the normal vaginal flora after antibiotic treatment of bacterial vaginosis. *BJOG.* 2008;115(11):1369-74.
124. Rossi A, Rossi T, et al. The use of *Lactobacillus rhamnosus* in the therapy of bacterial vaginosis. Evaluation of clinical efficacy in a population of 40 women treated for 24 months. *Arch Gynecol Obstet.* 2010;281(6):1065-9.
125. Larsson P, Brändborg E, et al. Extended antimicrobial treatment of bacterial vaginosis combined with human *Lactobacilli* to find the best treatment and minimize the risk of relapses. *BMC Infectious Diseases.* 2011;11:233.
126. Oduseyo O, et al. The effects of antimicrobial therapy on bacterial vaginosis in non-pregnant women. *Cochrane Collaboration.* 2009;1-77.
127. Martinez R, et al. Improved treatment of vulvovaginal candidiasis with fluconazole plus probiotic *Lactobacillus rhamnosus* GR-1 and *Lactobacillus reuteri* RC-14. *Letters in Applied Microbiology* 48 (2009) 269–274.
128. Martinez R, et al. Improved cure of bacterial vaginosis with single dose of tinidazole (2g), *L. reuteri* RC-14, and *L. rhamnosus* GR-1: a randomized, double-blind, placebo-controlled study. *Can J Microbiol.* 55:153-158, 2009;
129. Vujic G, et al. Efficacy of orally applied probiotic capsules for bacterial vaginosis and other vaginal infections: a double-blind, randomized, placebo-controlled study. *Eur J Obs&Gyn Rep Biol* 168 (2013) 75-79
130. Agrawal A, Houghton LA, et al. Clinical trial: The effects of a fermented milk product containing *Bifidobacterium lactis* DN-173 010 on abdominal distension and gastrointestinal transit in irritable bowel syndrome with constipation. *Alimenta*
131. Guyonnet D, Chassany O, et al. Effect of a fermented milk containing *Bifidobacterium animalis* DN-173 010 on the health-related quality of life and symptoms in irritable bowel syndrome in adults in primary care: A multicenter, randomized, dou
132. Yang WX, Mei H, et al. Effect of a fermented milk containing *Bifidobacterium lactis* DN-173010 on Chinese constipated women. *World J Gastroenterol.* 2008;14(40):6237-43.
133. Merenstein D, Murphy M, et al. Use of a fermented dairy probiotic drink containing *Lactobacillus casei* (DN-114 001) to decrease the rate of illness in kids: The DRINK study. A patient-oriented, double-blind, cluster-randomized, placebo-contr
134. Pedone CA, Bernabeu AO, et al. The effect of supplementation with milk fermented by *Lactobacillus casei* (strain DN-114 001) on acute diarrhoea in children attending day care centres. *Int J Clin Pract.* 1999;53(3):179-84.
135. Pedone CA, Arnaud CC, et al. Multicentre study of the effect of milk fermented by *Lactobacillus casei* on the incidence of diarrhea. *Int J Clin Pract.* 2000;54(9):568-71.
136. Sýkora J, Valecková K, et al. Effects of a specially designed fermented milk product containing probiotic *Lactobacillus casei* DN-114 001 and the eradication of *H. pylori* in children: A prospective randomized double-blind study. *J Clin Gastro*
137. Lin JS, Chiu YH, et al. Different effects of probiotic species/strains on infections in preschool children: a double-blind, randomized, controlled study. *Vaccine* 2009;27(7):1073-9.
138. Hickson M, D'Souza AL, et al. Use of probiotic *Lactobacillus* preparation to prevent diarrhoea associated with antibiotics: Randomised double blind placebo controlled trial. *BMJ.* 2007;335(7610):80.
139. Giovannini M, Agostoni C, Riva E, Salvini F, Ruscito A, Zuccotti GV, Radaelli G. A randomized prospective double blind controlled trial on effects of long-term consumption of fermented milk containing *Lactobacillus casei* in pre-school children with allergic asthma and/or rhinitis. *Pediatr Res.* 2007 Aug;62(2):215-20.
140. Merenstein D, Murphy M, Fokar A, Hernandez RK, Park H, Nsouli H, Sanders ME, Davis BA, Niborski V, Tondu F, Shara NM, Use of a fermented dairy probiotic drink containing *Lactobacillus casei* (DN-114 001) to decrease the rate of illness in kids: the DRINK study. A patient-oriented, double-blind, cluster-randomized, placebo-controlled, clinical trial. *Eur J Clin Nutr.* 19 Mai 2010 "on-line"
141. Boge T, Rémy M, Vaudaine S, Tanguy J, Bourdet-Sicard R, van der Werf S. A probiotic fermented dairy drink improves antibody response to influenza vaccination in the elderly in two randomised controlled trials. *Vaccine.* 2009 Sep 18;27(41):5677-84.
142. Guillemand E, Tondu F, Lacoin F, Schrezenmeir J. Consumption of a fermented dairy product containing the probiotic *Lactobacillus casei* DN-114 001 reduces the duration of respiratory infections in the elderly in a randomised controlled trial. *Br J Nutr.* 2009
143. Guillemand E, Tanguy J, Flavigny A, de la Motte S, Schrezenmeir J. Effects of consumption of a fermented dairy product containing the probiotic *Lactobacillus casei* DN-114 001 on common respiratory and gastrointestinal infections in shift workers in a randomized controlled trial. *J. Am. Coll. Nutr.* 2010 29:455-468
144. Marcos A, Wärnberg J, Nova E, Gómez S, Alvarez A, Alvarez R, Mateos JA, Cobo JM. The effect of milk fermented by yogurt cultures plus *Lactobacillus casei* DN-114001 on the immune response of subjects under academic examination stress. *Eur J Nutr.* 2004 Dec;43(6):381-9
145. Meyer AL, Micksche M, Herbecat I, Elmadaf I. Daily intake of probiotic as well as conventional yogurt has a stimulating effect on cellular immunity in young healthy women. *Ann Nutr Metab.* 2006;50(3):282-9.
146. Ortiz-Andreluchi A, Sánchez-Villalaga A, Rodríguez-Gallego C, Lemes A, Molero T, Soria A, Peña-Quintana L, Santana M, Ramírez O, García J, Cabrera F, Cobo J, Serra-Majem L. Immunomodulatory effects of the intake of fermented milk with *Lactobacillus casei* DN114001 in lactating mothers and their children. *Br J Nutr.* 2008 Oct;100(4):834-45
147. Tiollier E, Chennaoui M, Gomez-Merino D, Drogou C, Filaire E, Guezenne C. Effect of a probiotics supplementation on respiratory infections and immune and hormonal parameters during intense military training. *Mil Med.* 2007 Sep;172(9):1006-11.
148. Turchet P, Laurenzano M, Auboiron S, Antoine JM. Effect of fermented milk containing the probiotic *Lactobacillus casei* DN-114001 on winter infections in free-living elderly subjects: a randomised, controlled pilot study. *J Nutr Health Aging.* 2003;7(2):75-7.
149. Saavedra JM, Abi-Hanna A, et al. Long-term consumption of infant formulas containing live probiotic bacteria: tolerance and safety-1. *Am J Clin Nutr.* 2004;79:261-7
150. Sheu B, Wu JJ, et al. Impact of supplement with *Lactobacillus*-and *Bifidobacterium*-containing yogurt on triple therapy for *Helicobacter pylori* eradication. *Alimentary Pharmacology & Therapeutics.* 2002;16:1669-75.
151. Sheu B, Cheng H, et al. Pretreatment with *Lactobacillus*-and *Bifidobacterium*-containing yogurt can improve the efficacy if quadruple therapy in eradicating residual *Helicobacter pylori* infection after failed triple therapy. *Am J Clin Nutr.* 20
152. Wang KY, Li SN, et al. Effects of ingesting *Lactobacillus*-and *Bifidobacterium*-containing yogurt in subjects with colonized *Helicobacter pylori*. *Am J Clin Nutr.* 2004;80(3):737-41.
153. de Vrese M, Kristen H, et al. Probiotic *Lactobacilli* and *bifidobacteria* in a fermented milk product with added fruit preparation reduce antibiotic associated diarrhea and *Helicobacter pylori* activity. *J Dairy Res* 2011; 78:396-403.
154. Uchida, K, et al. Effect of fermented milk containing *Bifidobacterium lactis* BB-12® on stool frequency, defecation, fecal microbiota and safety of excessive ingestion in healthy female students. *J. Nutr. Food* 2005, 8, 39-51
155. Nishida, S, et al Effect of yogurt containing *Bifidobacterium lactis* BB-12 on improvement of defecation and fecal microflora of healthy female adults. *Milk Sci.* 2004, 53, 71-80.
156. Ejtahed H, et al Probiotic yogurt improves antioxidant status in type 2 diabetic patients. *Nutrition.* 2012 May;28(5):539-43.

THIS EDUCATIONAL TOOL HAS BEEN MADE POSSIBLE THROUGH
AN UNRESTRICTED EDUCATIONAL GRANT BY



Alliance for
Education on
Probiotics

ALLIANCE MEMBERS



Seroyal

For more information on the Alliance visit AEProbio.ca
For an online version of this guide visit probioticchart.ca

2016 version (document annually updated)